

2018 Quota Submissions for Commercially Harvested Macropods in Queensland

Prepared by: Macropod Management Program, Environmental Services and Regulation, Department of Environment and Science

© State of Queensland, 2018.

The Queensland Government supports and encourages the dissemination and exchange of its information. The copyright in this publication is licensed under a Creative Commons Attribution 3.0 Australia (CC BY) licence.



Under this licence you are free, without having to seek our permission, to use this publication in accordance with the licence terms.

You must keep intact the copyright notice and attribute the State of Queensland as the source of the publication.

For more information on this licence, visit <http://creativecommons.org/licenses/by/3.0/au/deed.en>

January 2018

Executive summary

The commercial macropod harvest in Queensland is focused on three species (red kangaroo *Macropus rufus*, eastern grey kangaroo *Macropus giganteus*, and common wallaroo *Macropus robustus*) located in four zones: no harvest zone, eastern harvest zone, central harvest zone and western harvest zone.

The harvest is administered through a quota and a quota submission which is released annually and outlines proposed quotas for each species in each zone for the following calendar year. Quotas are set between 10% and 20% of the estimated population for each species in each zone, depending on survey intensity and the standard error associated with population estimates.

Population estimates for 2017 and proposed sustainable use quotas for the 2018 commercial harvest

Species	Harvest zone	2017 estimated population (rounded to the nearest 50)	2018 sustainable use quota (rounded to the nearest 50)	Proportion of population (% rounded to the nearest whole number)
Red kangaroo	Central	5,866,450	1,173,300	20
	Eastern	174,400	17,450	10
	Western	485,850	48,600	10
	Combined	6,526,700	1,239,350	19
Eastern grey kangaroo	Central	8,681,600	1,213,050	14
	Eastern	4,303,300	430,350	10
	Western	7,250	0	0
	Combined	12,992,150	1,643,400	13
Common wallaroo	Central	1,790,200	268,550	15
	Eastern	1,579,850	158,000	10
	Western	47,050	4,700	10
	Combined	3,417,100	431,250	13

The formation of quotas is informed by criteria including:

- population trends (estimates obtained through aerial surveys)
- review of previous harvests
- the extent of non-commercial harvest mortality
- the extent of areas not subject to any harvest
- rainfall trends.

This quota submission outlines in detail the aspects of these criteria that pertain to the 2018 proposed quotas.

For 2017, aerial surveys were conducted at 13 monitor blocks across Queensland. Since regionalisation of the Queensland commercial macropod harvest was introduced in 2003, an estimate of macropod population size in the eastern and western zones has been made. The model used to estimate these populations is based on a small sample area and the reduced sampling effort is reflected in a conservative quota. This model was updated in 2012 to incorporate almost a decade of survey data and to generate trigger points for the commercial quota allocation.

Population estimates have decreased marginally for eastern grey kangaroos in the eastern zone but increased overall in the central zone in 2017. However in the central zone north the eastern grey population estimate is below a predetermined trigger point and the quota for this region has been halved for 2018. The red kangaroo population estimate for the central and eastern zones increased slightly since the 2016 surveys. Common wallaroo population estimates decreased in the central zone but increased significantly in the eastern zone. The western zone was not surveyed in 2017. Overall combined totals for all three species remained similar to 2016 across the state.

Examination of long-term trends in population and block density estimates indicates that the 2017 estimates are within the realm of fluctuations in previous years. Estimates for all three commercially harvested species consistently number over 1,000,000 in Queensland.

In the 2016 harvest period, 22.9% of the commercial harvest quota was utilised, with the highest percentage of quota used being 31.2% for eastern grey kangaroos in the central zone. The overall harvest was male biased, with females comprising less than 4% of the overall harvest.

Figures available on 31 May 2017 show that 5.3% of the available quota for red kangaroos in the western zone had been harvested with 3.3% and 10.2% of the quota harvested in the eastern and central zones respectively. For eastern grey kangaroos, 14.7% and 2.5% of the quota was harvested in the central and eastern zones respectively. For common wallaroos, the highest percentage of quota harvested was 17.9% in the central zone, whilst only 2% of the quota was harvested in the eastern zone and 8.7% in the western zone. Given these figures, it is unlikely that quotas will be met for each species in each zone in 2017.

Non-commercial take under damage mitigation permits (DMPs) were below quota for the 2016 harvest period. This trend is likely to be repeated in 2017 with a decrease in the uptake of DMPs by landholders.

The three commercially harvested macropod species continue to be protected from harvesting within the harvest zones in national parks and state forests. The protected area within the harvest zones is 79,981km². Macropods are further protected from harvest in Queensland within the non-harvest zones.

Annual rainfall across Queensland was wetter than average during 2016 with many parts of the state flooding in winter that year. However 2017 has been very dry throughout most parts with the majority of the harvest zones still drought declared at 31 August 2017.

Contents

Executive summary	iii
Introduction	1
Proposed quotas.....	2
Criteria considered in quota determination	4
Long-term population trends	4
Trigger points	11
Comparison between 2015 and 2016 population estimates.....	13
Review of the 2015 harvest.....	20
Sex ratio by species and zone	21
Harvest update for 2016.....	23
The extent of non-commercial harvest mortality	23
Proportion of the population not subject to harvesting.....	26
Rainfall trends	30
Summary and conclusion.....	33
References.....	34
Appendixes	35
Appendix 1 Summary of the methodology for population monitoring and quota derivation for Queensland	35
Appendix 2 Densities per km ² of the commercially harvested macropod species 2004–2016	38

Introduction

The commercial harvesting in Queensland of three macropod species—red kangaroo *Macropus rufus*, eastern grey kangaroo *Macropus giganteus* and common wallaroo *Macropus robustus*—is regulated through:

- *Environment Protection and Biodiversity Conservation Act 1999*
- Queensland Wildlife Trade Management Plan for Export—Commercially Harvested Macropods 2013–2017
- *Nature Conservation Act 1992*
 - Nature Conservation (Administration) Regulation 2017
 - Nature Conservation (Wildlife Management) Regulation 2006
 - Nature Conservation (Wildlife) Regulation 2006
 - Nature Conservation (Macropod) Conservation Plan 2017
 - Nature Conservation (Macropod Harvest Period) Notice
- *Animal Care and Protection Act 2001*
- *Food Production (Safety) Act 2000*.

The Department of Environment and Science (DES) administers the harvest of macropods in Queensland in accordance with the International Union for Conservation of Nature (IUCN) Recommendation 18.24, ‘the ethical, wise and sustainable use of some wildlife can provide an alternative or supplementary means of productive land-use, and can be consistent with and encourage conservation, where such use is in accordance with appropriate safeguards’ (IUCN 1990) and the Wildlife Trade Management Plan for Export—Commercially Harvested Macropods 2013–17, ‘to provide for the sustainable use of macropod species covered by the plan, in accordance with the principles of ecologically sustainable development’ (Anon 2012).

Management of the harvest is facilitated via a quota that sets the number of animals that can be taken. Quotas are established largely based on aerial surveys of the commercially harvested species and have been used in Queensland since 1975. The Director-General of DES declares a harvest period open annually via the harvest period notice and sets quotas for this period having regard to the Queensland Wildlife Trade Management Plan for Export—Commercially Harvested Macropods 2013–17. Quotas are provided to the Commonwealth Minister for The Environment for endorsement.

Quotas in Queensland are set between 10% and 20% of the estimated population for each species in each zone. Harvesting at these levels will ensure a sustainable yield and long-term conservation of macropod populations.

Since 2003, quotas have been set for each species for four harvest zones to ensure that harvest pressure is distributed across the range of the species (Figure 1):

- no harvest zone (quota zero)
- eastern harvest zone
- central harvest zone
- western harvest zone.

This quota submission contains a summary of the recommended quotas for each of the species in each of the harvest zones for 2018. Additionally, the submission outlines the basis of how these quotas were determined.

The Nature Conservation (Macropod Harvest Period 2018) Notice 2017 is due for release in December 2017. The release of this notice will allow the harvest period to be declared open on 1 January 2018. The notice will outline specific conditions for the 2018 harvest period including, but not limited to, harvest zone boundaries, weights and more.

The proposed quotas were calculated using a fixed proportion of the estimated macropod populations within the Queensland harvest areas. Proportions were adjusted for each species across the harvest zones in relation to the margins of error present in population estimates derived from aerial surveys. The maximum proportions used for each species were 15% of the populations for eastern grey kangaroos and common wallaroos and 20% of the population for red kangaroos for the central zone. For the eastern and western zones, where survey effort is less extensive when compared to the central zone, the more conservative maximum proportion of 10% was applied for all three species.


These sustainable-use harvest proportions are based on research and modelling undertaken by Caughley et al. (1987) and Hacker et al. (2002) and are currently accepted by the scientific community, DES and the Department of the Environment and Energy for determining state quota limits.

Proposed quotas

Table 1 – 2017 estimated populations and 2018 proposed quotas for each macropod species in each harvest zone

Species	Harvest zone	2017 estimated population	2018 sustainable use quota (rounded to the nearest 50)	Proportion of population (% rounded to the nearest whole number)
Red kangaroo	Central	5,866,450	1,173,300	20
	Eastern	174,400	17,450	10
	Western	485,850	48,600	10
	Combined	6,526,700	1,239,350	19
Eastern grey kangaroo	Central	8,681,600	1,213,050	14
	Eastern	4,303,300	430,350	10
	Western	7,250	0	0
	Combined	12,992,150	1,643,400	13
Common wallaroo	Central	1,790,200	268,550	15
	Eastern	1,579,850	158,000	10
	Western	47,050	4,700	10
	Combined	3,417,100	431,250	13

Figure 1 – Queensland macropod harvest zones



Criteria considered in quota determination

The following criteria were considered in determining the quotas for each macropod species:

- population trends (estimates obtained through aerial surveys)
- review of previous harvests
- the extent of non-commercial harvest mortality
- the extent of areas not subject to any harvest
- rainfall trends.

Each of these criteria will be outlined in detail in this quota submission.

Long-term population trends

Population estimates

Since 1992, the Queensland Government has coordinated an annual program of aerial surveys to directly monitor populations of the three commercially harvested macropod species. These surveys occur over 22 representative monitor blocks across the state and are utilised to obtain population estimates that inform the quota. The methodology of the surveys is outlined in detail in Appendix 1. Since 2011 a correction factor of 1.85 has been applied to population estimates for common wallaroo in Queensland. Prior to this a correction factor of 1.2 was applied.

Current harvesting rates (quotas ranging from 10% to 20% of population estimates) are considered sustainable. None of the three commercially harvested species has shown a consistent decline in abundance since 1992 (figures 2, 3, 4 and 5), which would necessitate a reassessment of the harvest take and species conservation status. Whilst no consistent declines have been observed, the macropod populations in Queensland have fluctuated over time.

Density estimates

To contribute to ensuring commercially harvested macropod species are maintained across their distributions, density estimates are calculated for representative survey blocks as part of the aerial surveys (Appendix 2). For the purposes of interpreting this data in an easily understood manner, the density estimates for each species have been grouped into the eastern and western harvest zones, with the central harvest zone divided into three regions, central zone north (Mitchell Grass dominant), central zone south (Mulga Lands dominant) and central zone east (Brigalow Belt dominant) (Figure 6).

Figure 2 – Macropod population trends—1992–2017




Figure 3 – Red kangaroo population estimates (with one standard error) since 1992




Figure 4 – Eastern grey kangaroo population estimates (with one standard error) since 1992





Figure 5 – Common wallaroo population estimates (with one standard error) since 1992. Estimates include a 1.85 correction factor for 2011 to 2017 and a 1.2 correction factor all other years



This data is monitored for any significant decreases in densities. This approach is possible for all data collected since 2005. Examination of trends in density for the three commercially harvested macropod species in the areas outlined above for the period 2005–2017 demonstrates densities do fluctuate over time (figures 7–11).

For red kangaroos densities are greatest in the central zone north and central zone south, with densities lowest in the eastern zone (Figure 10). Low densities in the eastern zone are expected as this area incorporates the edge of the distributional range for this species.

For eastern grey kangaroos highest densities are recorded in the central zone east (Figure 9). Eastern grey kangaroos occur in consistently low densities in the western zone at the edge of their distributional range. As such, there is no quota for eastern grey kangaroos in this zone (Table 1).

Common wallaroos occur in highest densities in the central zone north. This area is further divided into two regions for the purposes of estimating populations due to the considerably higher densities recorded around Blackall, Tambo and Longreach when compared to the rest of the zone (Figure 7 and Appendix 2). Lowest densities for this species occur in the western zone (Figure 11). Density fluctuations for this species do not follow the same patterns as those exhibited by red and eastern grey kangaroos (figures 7–11).

Figure 6 – Regions used to calculate population estimates of commercially harvested macropods




Figure 7 – Average density km² of commercially harvested macropods in the Central North population estimate region from 2005 to 2017 (Common wallaroos are represented by two areas in the Central North zone).




Figure 8 – Average density km² of commercially harvested macropods in the Central South population estimate region from 2005 to 2017




Figure 9 – Average density km² of commercially harvested macropods in the Central East population estimate region from 2005 to 2017




Figure 10 – Average density km² of commercially harvested macropods in the Eastern population estimate region from 2005 to 2017





Figure 11 – Average density km² of commercially harvested macropods in the Western population estimate region from 2005 to 2017



Trigger points

2013 marked the beginning of a new Queensland Wildlife Trade Management Plan for Export (Commercially Harvested Macropods 2013–17). Incorporated into this new plan are pre-determined trigger points for each of the commercial harvest quotas. Each trigger point represents a threshold level based on analysis of the long-term population estimate for each harvested species in each population estimate region.

Where an estimated population for a region falls below a trigger point of 1.5 standard deviations (SD) below the long-term average for that region then the harvest quota will be halved for that region in the next calendar year. If a population estimate falls below 2 SDs below the long-term average for that species in that region then there will be no quota for the following year.

Table 2 shows the calculated trigger points for 2018 for each species in each region compared with the population estimates for those regions. The estimated population for eastern grey kangaroos in the central north region is below 1.5 SD trigger point. Consistent with the Wildlife Trade Management Plan for Export—Commercially Harvested Macropods 2013–17 the harvest quota for this species has been halved for 2018. The harvest of eastern grey kangaroos will be limited to a maximum of 89,200 in the central north region for 2018 (figure 12). All other species are above the calculated trigger points and quotas have been allocated accordingly (Figure 12). As the western zone is on the very limit of the eastern grey kangaroo range no quota is set for this species in this zone. Consequently there are no trigger points calculated for this species in this zone.

Table 2 – Calculated trigger points for 2018 and estimated populations of commercially harvested macropod species in each region for 2017

Note: There is no quota set for eastern grey kangaroos in the western region. The quota for this species will be halved for the central north region in 2018 as the estimated population is below the 1.5 SD trigger point.

Species	Population estimate region	2017 estimated population	2018 1.5 SD trigger point	2018 2 SD trigger point
Red kangaroo	Central North	3,384,364	1,815,176	1,599,711
	Central South	1,949,256	689,013	557,976
	Central East	532,808	96,044	72,409
	Eastern	174,408	73,800	63,800
	Western	485,835	180,150	142,400
Eastern grey kangaroo	Central North	1,189,223	1,236,292	1,017,324
	Central South	1,560,788	861,140	692,537
	Central East	5,931,583	2,450,768	2,068,980
	Eastern	4,303,285	1,152,100	886,300
	Western	7,267	NA	NA
Common wallaroo	Central North	1,185,629	444,645	338,405
	Central South	504,748	93,187	67,614
	Central East	99,803	15,587	11,860
	Eastern	1,579,844	209,650	164,700
	Western	47,057	20,300	12,300

Note: There is no quota set for eastern grey kangaroos in the western region.

Whilst the effect of halving the harvest quota for eastern greys in the central north reduces the available harvest for this species this effect is unlikely to impact the industry. The overall availability of harvest macropods in the central north is 943,900 whilst the overall availability of eastern greys in the entire central zone is 1,213,050 (figure 12). Although the estimated population for eastern greys decreased in this region it remained very similar to 2016 and increased slightly over the combined central harvest zone (figure 13).

Figure 12- Detail of the calculated harvest quotas for the central zone including the reduction of eastern grey kangaroos in the central north.





Figure 13- Comparison of eastern grey kangaroo estimated populations between 2016 and 2017 in the central harvest zone.



Comparison between 2016 and 2017 population estimates

The total population estimates combined across all three harvest zones has decreased for eastern grey kangaroos and increased for red kangaroos and common wallaroos in 2017 compared with 2016 (Figure 13). However whilst some population estimates have decreased for specific zones others have increased (Table 3). Red kangaroo population estimates increased in the central and eastern zones (figure 14). The eastern grey population estimates decreased in the eastern zones and increased in the central zone (figure 14). Common wallaroo population estimates increased significantly in the eastern zone due mostly to a large density increase recorded at the Inglewood survey block (Appendix 2). However common wallaroos decreased in the central zone (figure 14). The western harvest zone was not surveyed in 2017.

Since regionalisation of the Queensland commercial macropod harvest was introduced in 2003 an estimate of macropod population size in the eastern and western zones has been made. The model used to estimate these populations is based on a small sample area and the reduced sampling effort is reflected in a conservative quota (figure 15). This model was updated in 2012 to incorporate almost a decade of survey data and to generate trigger points for the commercial quota allocation.

Figure 13 – Comparison of overall macropod populations in the commercial harvest zones 2016 and 2017 (with one standard error).




Table 3 – Comparison between 2016 and 2017 macropod population estimates. Note the western zone was not surveyed in 2017.

Species	Harvest zone	2016 population estimate	2017 population estimate
Red kangaroo	Central	5,137,600	5,866,450
	Eastern	145,200	174,400
	Western	485,850	485,850
	Combined	5,768,650	6,526,700
Eastern grey kangaroo	Central	8,111,850	8,681,600
	Eastern	5,873,850	4,303,300
	Western	7,250	7,250
	Combined	13,992,950	12,992,150
Common wallaroo	Central	2,205,600	1,790,200
	Eastern	552,500	1,579,850
	Western	47,050	47,050
	Combined	2,805,150	3,417,100

Figure 14 – Comparison of macropod populations from 2016 to 2017 by species and zone.






Figure 15 – Comparison of 2017 actual and 2018 proposed harvest quotas.



Long-term quota and harvest trends


Figures 16 to 18 outline data on the three commercially harvested macropod species pertaining to estimated population, quota and harvest for the years 1992 to 2016. Please note that population estimates are based on aerial surveys conducted in the previous year to the quota and harvest. Combined population estimates, quota and harvest data have been used for the period post-regionalisation in 2003, to enable comparison with data collated prior to this period.

Figure 16 – Long-term population estimates (\pm one standard error), quota and harvest data (commercial harvest + damage mitigation permits) for the red kangaroo in Queensland.




Note: population estimates are based on aerial surveys conducted the year before the harvest was taken.

Figure 17 – Long-term population estimates (\pm one standard error), quota and harvest data (commercial harvest + damage mitigation permits) for the eastern grey kangaroo in Queensland.



Note: population estimates are based on aerial surveys conducted the year before the harvest was taken.

Figure 18 – Long-term population estimates (\pm one standard error), quota and harvest data (commercial harvest + damage mitigation permits) for the common wallaroo in Queensland.



Note: population estimates are based on aerial surveys conducted the year before the harvest was taken.

As previously outlined, there has been no consistent decline in the populations of the three commercially harvested species since 1992 (figures 16 to 18). Of these species, the eastern grey kangaroo is consistently most abundant across the harvest zones, followed by the red kangaroo. Common wallaroos are the least numerous. Population estimates for all three species are in excess of one million across the harvest zones.

As quotas are set as a constant proportion of the populations, they fluctuate as population estimates fluctuate (figures 16 to 18). However, numerous factors influence harvest rates for commercial macropods. These include population levels, market forces, environmental conditions and access by harvesters. As a consequence, there is no clear pattern or trend in the proportion of the quota harvested since 1992.

Review of the 2016 harvest

Dealer returns for the year 2016 indicate that there were 896,144 macropods taken in Queensland, which represents 22.9% of the overall combined quota. Of the 896,144 animals harvested, there were 251,778 red kangaroos, 528,547 eastern grey kangaroos and 115,819 common wallaroos harvested (table 4). Quotas for individual species in each harvest zone were not exceeded in 2016, the maximum commercial take as a percentage of the approved quota of 31.2% being for eastern greys in the central zone (tables 4 to 7).

Table 4 – Total harvest in 2016

Species	Population estimate 2015	Quota 2016	Harvest take 2016	% quota used 2016	% population harvested 2016
Red kangaroo	7,061,200	1,335,950	251,778	18.8%	3.6%
Eastern grey kangaroo	15,406,700	2,055,250	528,547	25.7%	3.4%
Common wallaroo	3,694,100	525,500	115,819	22.0%	3.1%
Total	26,162,000	3,916,700	896,144	22.9%	3.4%

Note: population estimates are based on aerial surveys conducted in 2015, which were used to set the 2016 quota.

Table 5 – Harvest of red kangaroos in 2016

Zone	Population estimate 2015	Quota 2016	Harvest take 2016	% quota utilised 2016	% population harvested 2016
Central	6,298,150	1,259,650	240,857	19.1%	3.8%
Eastern	171,500	17,150	2,360	13.8%	1.4%
Western	591,550	59,150	8,561	14.5%	1.4%
Total	7,061,200	1,335,950	251,778	18.8%	3.6%

Note: population estimates are based on aerial surveys conducted in 2015, which were used to set the 2016 quota.

Table 6 – Harvest of eastern grey kangaroos in 2016

Zone	Population estimate 2015	Quota 2016	Harvest take 2016	% quota utilised 2016	% population harvested 2016
Central	10,291,600	1,543,750	481,666	31.2%	4.7%
Eastern	5,115,100	511,500	46,881	9.2%	0.9%
Western	0	0	-	-	0.0%
Total	15,406,700	2,055,250	528,547	25.7%	3.4%

Note: population estimates are based on aerial surveys conducted in 2015, which were used to set the 2016 quota.

Table 7 – Harvest of common wallaroos in 2016

Zone	Population estimate 2015	Quota 2016	Harvest take 2016	% quota utilised 2016	% population harvested 2016
Central	3,130,200	469,550	110,699	23.6%	3.5%
Eastern	559,400	55,950	5,120	9.2%	0.9%
Western	4,500	0	-	-	-
Total	3,694,100	525,500	115,819	22.0%	3.1%

Note: population estimates are based on aerial surveys conducted in 2015, which were used to set the 2016 quota.

Sex ratio by species and zone

Commercial harvest is typically biased towards males (figure 20) as they are usually larger and heavier than females. In 2016, the total harvest for each species comprised between 95.7% and 99.9% males. Data gathered throughout 2016 indicates 3.4% of the overall harvest was female (figure 19).

Figure 19 - Proportion of 2016 harvest, male and female by species





Figure 20 – Overall sex ratio from 2006 to 2016 (percentages rounded to the nearest whole number).



Harvest update for 2017

The total number of tags issued up to 31 May 2017 was 683,400. A comparison of tag sales and harvest returns in relation to quotas in each zone is given in Table 8. The number of tags produced does not exceed the maximum quota for each species in each zone therefore it is not possible for the commercial harvest quotas to be over allocated. Given the low percentage of the quota that has currently been issued in tags, it is unlikely quotas will be met for any species in any zone. The 2017 harvest will be comprehensively reported on in the 2017 annual report, due for release in March 2018.

Table 8 – Tags issued and reported harvest for 2017 at 31 May.

Species	Harvest zone	2017 sustainable use quota (rounded to the nearest 50)	Tags issued to 31 May 2017	Reported harvest to 31 May 2017
Red kangaroo	Central	1,027,500	167,850	104,756
	Eastern	14,500	2,600	483
	Western	48,600	5,550	2,588
Eastern grey kangaroo	Central	1,216,800	272,400	179,049
	Eastern	587,400	31,800	14,578
	Western	NA	NA	NA
Common wallaroo	Central	330,850	95,150	59,150
	Eastern	55,250	5,200	1,120
	Western	4,700	1,050	410


The extent of non-commercial harvest mortality

There are many forms of macropod mortality outside of the commercial harvest. It is possible for DES to collect and report data on three forms of non-commercial harvest mortality which can be considered when determining commercial quotas. These include Damage Mitigation Permits (DMPs), licensed recreational harvest and disease outbreak mortality.

Damage to primary production


DMPs are issued by DES where macropods are causing demonstrable damage to primary production. The issuing of these permits is limited to a maximum of 2% of the population estimate for each species in each zone. It is a condition of the permit that macropods are taken in accordance with the requirements of the National Code of Practice for the Humane Shooting of Kangaroos and Wallabies for Non-commercial Purposes. Uptake of DMPs is below the available quota (figure 21) and has decreased since 2015 (figure 22).

Figure 21 – Macropod quota and take for DMP in 2017.



Note: Figures are as recorded on 14 September 2017

Figure 22 – Macropods taken under a DMP 2006–2016.



Disease outbreak mortality and its significance

Following local media reports from Longreach in October 2016 Biosecurity Queensland sent a team to Noonbah station to investigate potential macropod deaths. After 2 days of extensive searching no evidence was found of any recent macropod mortalities. No incidence of significant disease mortalities have been recorded for macropod populations in Queensland during 2017. Whilst the winter of 2016 was wetter than average above average temperatures, extremely dry conditions, have continued across majority of the commercial harvest zones in 2017. Most of the local government areas within the harvest zones have been drought declared for over three years. These conditions reduce the available feed for all grazing species including macropods. The main areas to experience a significant reduction in macropod numbers is in the west of the state and for common wallaroos the central north.

The last disease related non-harvest mortality event recorded in Queensland macropod populations was in March 2010. Localised disease related mortalities for some commercially harvested species were recorded in the south-west of the state from the New South Wales border to just north of Quilpie in the Paroo and Bulloo river systems. Investigation and surveillance over the weeks following the reports established that the deaths were very limited and restricted to small areas only. This was documented in the 2011 Quota Submission for Queensland. The aerial surveys conducted in 2011 showed that in the year following the reported mortalities, macropod populations had increased indicating that the reported mortalities in 2010 did not have a significant effect on macropod populations in the area.

Licensed recreational wildlife harvesting licence (macropods)

The recreational macropod harvest in Queensland is centred on the same three species as the commercial harvest and runs for the same harvest period. The recreational harvesting of these macropods is regulated through:

- *Environment Protection and Biodiversity Conservation Act 1999*
- *Nature Conservation Act 1992*
 - Nature Conservation (Administration) Regulation 2006 (current to September 2017)
 - Nature Conservation (Administration) Regulation 2017 (current from September 2017)
 - Nature Conservation (Wildlife Management) Regulation 2006
 - Nature Conservation (Wildlife) Regulation 2006
 - Nature Conservation (Macropod) Conservation Plan 2005 (current to September 2017)
 - Nature Conservation (Macropod) Conservation Plan 2017 (current from September 2017)
 - Nature Conservation (Macropod Harvest Period) Notice
- *Animal Care and Protection Act 2001.*

A person holding a Commercial Wildlife Harvesting Licence (macropods) cannot hold a Recreational Wildlife Harvesting Licence (macropods). The harvest is restricted to a maximum of 50 tags per licence holder during a harvest period. The recreational harvest of macropods in Queensland is small with a total of 1,049 tags issued in 2017 to 23 Licence holders (Recreational Wildlife Harvesting Licence) (figures current at 31 August 2017).

2017 is the last year Recreational Wildlife Harvesting Licences (macropods) will be issued as they have been combined with the Commercial Wildlife Harvesting Licence (macropods) in the Nature Conservation (Macropod) Conservation Plan 2017. From 2018 onwards a Macropod Harvesting Licence will be required to harvest macropods for commercial or personal use.

Proportion of the population not subject to harvesting

Commercial harvesting of macropods can only occur in three harvest zones in Queensland. Cape York Peninsula and the south east corner of Queensland are designated Non-harvest zones (figure 23). Within the three commercial harvest zones macropods cannot be harvested within National Parks, States Forests, Regional Parks, Timber Reserves and Forest Reserves. Table 9 outlines the size of these land tenures within the commercial harvest zones (current in 2017).

Figures 23 to 25 show the general distribution of each of the commercially harvested macropods in relation to the population estimate regions. Eastern grey kangaroos are only harvested in the eastern and western harvest zones (figure 24). Red kangaroos are harvested in the western and central harvest zones and in the north of the eastern harvest zone (figure 23). Common wallaroos have the broadest distribution (figure 25) throughout Queensland and can be harvested in the central, eastern and western zones.

Table 9 – Area of land tenures within the Queensland commercial harvest zones where harvesting of macropods is not permitted.

	Western harvest zone km ²	Central harvest zone km ²	Eastern harvest zone km ²	Total km ²
National Park	21,450	11,778	16,255	49,483
State Forest	NA	10,788	16,254	27,042
Regional Park	1,048	167	1,365	2,580
Forest Reserve	NA	NA	225	225
Timber Reserve	79	451	121	651
Total km ²	22,577	23,184	34,220	79,981

Figure 23 – Red kangaroo *Macropus rufus* distribution.




Figure 24 – Eastern grey kangaroo *Macropus giganteus* distribution.





Figure 25 – Common wallaroo (*Macropus robustus*) distribution.



Rainfall trends

Rainfall across western Queensland in 2016 was above average with the second wettest winter recorded since 1912 (figures 26 & 27). Extensive flooding occurred in some parts of the harvest zones. Temperatures were above average across the state throughout the year with April and May being the hottest ever recorded. The first half of 2017 has been warmer than average and dry for much of the harvest zones. Drought declarations continue across the majority of areas where macropod harvesting occurs (figure 28).

Much of the commercial harvest zones have now been drought declared for five consecutive years. Despite the ongoing drought conditions sufficient rainfall has occurred in some parts of the harvest zones over the last 12 months to produce feed for macropods. The population estimate for eastern grey kangaroos in the central north region was below the 1.5 SD trigger point in 2017. However the overall population estimates are similar to 2016 with no dramatic decline or increase observed in any of the harvest zones. Given the continued dry conditions the potential exists for macropod densities to decrease dramatically like they did in 2002 (figure 26) across the harvest zones.

Figure 26 Estimated macropod populations plotted with annual rainfall totals from Queensland from 1992 to 2016.




Figure 27 – Queensland rainfall totals (mm) from 1 September 2016 to 31 August 2017.





Figure 28 – Queensland drought declarations at 1 July 2017.



Summary and conclusion

The proposed quotas for the 2018 commercial macropod harvest in Queensland have been formulated by following an established methodology, which is largely based on constant proportions of population estimates and monitoring of long-term population trends. Population estimates are derived from representative aerial surveys across the harvest zones that are informed by the best available science. Other factors considered in the setting of the 2018 commercial harvest quotas include a review of previous harvests, extent of non-commercial harvest, proportion of the population not subject to harvesting, non-commercial harvest mortality and its significance, and rainfall trends.

Long-term trend data relating to population size since 1992, when Queensland began an annual program of helicopter surveys, demonstrates there has been no consistent increase or decline in the populations of red kangaroos, eastern grey kangaroos or common wallaroos in Queensland. However, populations do fluctuate over time. Population estimates for all species indicate that numbers over 1,000,000 occur in the harvest area. Thus, current harvest rates can be viewed as not having a long-term detrimental impact on populations.

Since regionalisation of the Queensland commercial macropod harvest was introduced in 2003, an estimate of macropod population size in the eastern and western zones has been made. The model used to estimate these populations is based on a small sample area and the reduced sampling effort is reflected in a conservative quota. This model was updated in 2012 to incorporate almost a decade of survey data and to generate trigger points for the commercial quota allocation. The population estimates in the eastern and western zones are a function of both the new model and the survey data for 2017.

Population estimates have decreased marginally for eastern grey kangaroos in the eastern zone but increased overall in the central zone in 2017. However in the central zone north the eastern grey population estimate is below a predetermined trigger point and the quota for this region has been halved for 2018. The red kangaroo population estimate for the central and eastern zones increased slightly since the 2016 surveys. Common wallaroo population estimates decreased in the central zone but increased significantly in the eastern zone. The western zone was not surveyed in 2017. Overall combined totals for all three species remained similar to 2016 across the state.

For the 2016 commercial harvest no quotas were exceeded, with the maximum percentage of quota utilised, being 31.2% for eastern kangaroos in the central zone. Sex ratios from harvest data continue to be biased towards males with the overall percentage of females harvested below 4%. Thus, the last completed harvest period provides no indication of adverse pressure on populations that would influence proposed quotas.

For the 2017 harvest period up to 31 May, 5.3% of the available quota for red kangaroos in the western zone had been harvested with 3.3% of the quota harvested in the eastern, and 10.2% in the central zones respectively. For eastern grey kangaroos, 14.7% of the quota was harvested in the central and 2.5% in the eastern zones respectively. For common wallaroos, the highest percentage of quota harvested was 17.9% in the central zone, with 2% harvested in the eastern zone and 8.7% in the western zone. Given these figures, it is unlikely that quotas will be met for each species in each zone in 2017.

Usage of DMPs in 2016 were below the 2% of the population estimate quota for all species for all zones. The current percentages for usage of DMP quotas for 2017 are below the new quota limit of 2%.

The three commercially harvested macropod species are protected from harvesting within the harvest area through national parks and state forests. These 'refuges' occur in patches throughout the distributional ranges of all three species. Macropods are further protected from harvest in Queensland within the non-harvest zones.

Overall numbers of commercially harvested macropods in Queensland harvest zones has remained relatively stable over the last 12 months. The majority of the harvest zones are drought declared and macropod numbers overall have decreased slightly. Should the widespread dry conditions continue throughout the state it is possible that observed macropod numbers will decrease again in 2018.

References

- Anon. 2012. Wildlife Trade Management Plan for Export—Commercially Harvested Macropods – 2013–2017. Queensland Parks and Wildlife Service; Brisbane.
- Anon. 2002. Macropod Commercial Quota Submission for 2002. Queensland Parks and Wildlife Service; Brisbane.
- Buckland, S.T., Anderson, D.R., Burnham, K.P., and J.L. Laake. 1993. Distance Sampling: Estimating Abundance of Biological Populations. Chapman and Hall; London.
- Cairns, S.C., Lollback, G.W., Payne, N. 2008. Design of aerial surveys for population estimation and the management of macropods in the Northern Tablelands of New South Wales, Australia. *Wildlife Research* 35: 331–339.
- Caughley, G., Shepard, N. and G. Short. 1987. Kangaroos, their ecology and management in the sheep rangelands of Australia. Cambridge University Press; Cambridge.
- Clancy, T.F., Edwards, G.P., Pople, A.R. and G.W. Maag. 1994. The 1993 surveys of the commercially harvested species of macropod in Queensland. Internal Report to Conservation Strategy Branch, Queensland Parks and Wildlife Service; Brisbane.
- Clancy, T.F., Pople, A.R., and L.A. Gibson. 1997. Comparison of helicopter line transects with walked line transects for estimating densities of kangaroos. *Wildlife Research* 24: 397–409.
- Hacker, R., McLeod, S., Druhan, J., Tenhumberg, B. and U. Pradhan. 2002. Managing Kangaroos in the Murray-Darling Basin. Technical Report to the Murray-Darling Basin Commission; Canberra.
- Pople, A.R., Cairns, S.C., Clancy, T.F., Grigg, G.C., Beard, L.A. and C.J. Southwell. 1998. Comparison of surveys of kangaroos in Queensland using helicopters and fixed-wing aircraft. *The Rangeland Journal* 20: 92–103.
- Pople, A.R. 2006. Modelling the spatial and temporal dynamics of kangaroo populations for harvest management. Final report to the Department of Environment and Heritage; Canberra.

Appendixes

Appendix 1 Summary of the methodology for population monitoring and quota derivation for Queensland

Aerial surveys

Since 1992, the Queensland Government has conducted an annual program of aerial surveys by helicopter to directly monitor populations of the three large macropod species covered by the Wildlife Trade Management Plan for Export—Commercially Harvested Macropods 2013–2017. This method employs line transect methodology (Buckland et al. 1993), which is significantly more robust to variations in sightability than standard fixed-wing methods and provides more accurate and precise population estimates (Clancy et al. 1997). A detailed description of the methodology employed in these surveys is provided in Clancy et al. (1997).

Surveys are conducted over 22 fixed monitor blocks, covering an area of 136,000km² (Figure 1) or 25% of the original fixed-wing survey area sampled by the Australian National Parks and Wildlife Service (ANPWS) between 1984 and 1995. In each helicopter survey block, between two to eight east-west running 50–90km transect lines have been placed systematically 10km apart. Sampling intensity within each block is approximately 2.5%.

The placement of the original 10 of the 22 survey blocks used in this monitoring program was designed to provide appropriate coverage of representative densities of macropods over the core harvest area of 630,000km² (Pople et al. 1998). In response to the introduction of regional management to Queensland in 2003, a further 12 survey blocks were added to provide broader coverage of the entire harvest area and to ensure all bioregions were sampled. Placement of these new survey blocks was optimised using fixed-wing survey data collected across the harvest zone during 2001. These surveys provided data to reassess the representativeness of the existing helicopter monitor blocks and to investigate alternative scenarios for future survey design. Analysis of these data in combination with harvest data has led to improved stratification of the survey area and hence increased the power to extrapolate data from survey monitor blocks to other regions in the harvest zone. These investigations were conducted in collaboration with the University of Queensland as part of the Australian Research Council (ARC) funded ROOSPIRT Linkage project (Pople et al. 2006).

No correction factors are applied to surveys of eastern grey and red kangaroos as comparisons of ground and aerial surveys conducted by Clancy et al. (1997) concluded that the helicopter line transect technique is both accurate and precise in determining population densities for both these species over a range of habitats, seasons and densities. Whilst the method is less accurate for common wallaroos there are still close correspondences between the results of helicopter surveys and those of ground counts. Estimates derived from ground surveys for common wallaroo density are approximately 1.9–2.0 times that recorded for helicopter surveys. Accordingly, since 1998, the Queensland Government has applied a conservative correction factor of 1.2 to the wallaroo density estimates derived from helicopter surveys. In 2011, this conservative correction factor has been increased to 1.85 in line with that used in New South Wales.

Frequency and coverage of aerial surveys

Due to the costs and logistics associated with conducting helicopter aerial surveys over 22 monitor blocks, not all monitor blocks are surveyed on an annual basis. To ensure adequate coverage of the harvest area in Queensland and to enable accurate tracking of population trends, all blocks are surveyed at least once every two years. Decisions on the frequency and coverage of the aerial survey program were based on analyses completed by the University of Queensland as part of the ARC funded ROOSPIRT Linkage project. This redesign of the monitoring program also sought to introduce a level of bioregional stratification to provide for improved potential to monitor populations at that scale as a possible prelude to the introduction of further harvest zones in Queensland.

The main feature of the current monitoring program is the establishment of pairs of closely correlated monitor blocks within each of the bioregions. The process of pairing monitor blocks was based on linear modelling that examined the relationships between macropod population densities, rainfall and harvest off-take for all monitoring blocks within each bioregion. The process of pairing blocks and the development and refinement of these linear models enables the frequency of monitoring of blocks to

be reduced without compromising the efficacy of the monitoring program. Using this approach monitoring blocks with bioregional pairs will be monitored on a rotating basis with each block being subject to surveys every two years.

Linear models utilising data on rainfall, harvest off-take and population rates of increase for the monitored block within a pair are then used to predict population changes in the unmonitored block of the pair. The models are refined annually as further data is collated and enhancements are made to the modelling process. In the situation where there is only a single monitor block within a bioregion or where a monitor block samples a unique macropod community, such as the Blackall block which contains moderate densities of all three species and particularly high densities of common wallaroo, these blocks are sampled annually. In order to calibrate the models and to provide a benchmark of the states' macropod populations, a survey of the complete set of 22 monitor blocks will be completed every five years. The table below summarises the current stratification, pairing and sampling frequency for all 22 of the fixed aerial survey monitor blocks used in the Queensland monitoring program.

Due to the significantly lower macropod densities and associated lower harvest densities recorded historically from the eastern and western harvest zones, these two zones are not surveyed with the same intensity as the central harvest zone. Within both the eastern and western harvest zones, population density estimates are based on helicopter aerial surveys of three monitor blocks established in 2004 and monitored on a two yearly rotation. The results are used in association with data derived from fixed-wing aerial surveys conducted over parts of these zones during 2001 to adjust approximations of the population rates of increase calculated by comparing fixed-wing survey data from 1981–82, 1984 and 2001 with the most recent helicopter aerial survey data and past ground surveys completed in equivalent habitats.

Population estimation

Population estimates are calculated by extrapolating the mean monitor block densities to a larger harvest area of 1,097,410km² for eastern grey kangaroos, 1,105,587km² for red kangaroos and 1,104,222km² for common wallaroos. To improve precision and remove bias in density estimates used in the calculation of population sizes and their standard errors, the data collected since 2003 have been stratified (Buckland et al. 1993) by observer and bioregion. Helicopter surveys are conducted with two observers, which results in twice the sampling intensity as one observer. To account for differences between observers, the data was post-sampling stratified by applying 'goodness of fit' models to the data from each observer using the computer program 'Distance' (Buckland et al. 1993). 'Distance' was then used to obtain an overall macropod density estimate for the survey block. The survey blocks were stratified by bioregion, and their weighted (by sampling intensity) density estimates used to calculate population sizes for the harvest area according to the methods and calculations outlined in Clancy et al. 1994 and 1997. For all species, stratification of the data by observer and bioregion has not significantly changed the population size estimate but has reduced the standard error associated with that estimate. However, with common wallaroos the standard errors associated with the population estimate remain high due to the high variability in densities across the state.

Quota derivation

Sustainable harvest quotas are calculated using a fixed proportion of the estimated macropod populations within the Queensland harvest area. The proportions used vary between species and are adjusted across the harvest zones in relation to the margins of error present in population estimates. The maximum proportions used for each species are 15% of populations for eastern grey kangaroos and common wallaroos and 20% of the population for red kangaroos. These maximum proportions are only applied to populations within the central harvest zone where survey effort is greatest and hence confidence limits for population estimates are within acceptable limits. In both the eastern and western harvest zones more conservative harvest proportions of 10% for all three species are applied. These sustainable-use harvest proportions are based on research and modelling undertaken by Caughley et al. (1987) and Hacker et al. (2002) and are currently accepted by the scientific community, the Queensland Department of Environment and Science and the Commonwealth Department of Sustainability, Environment, Water, Population and Communities for determining state quota limits.

Stratification, pairing and sampling frequency for fixed aerial survey monitor blocks in Queensland

Harvest zone	Bioregion	Monitor block	Years 1 and 3	Years 2 and 4	Year 5
Central	Brigalow Belt South	Injune	✓		✓
		Taroom		✓	✓
		Westmar	✓		✓
		Roma		✓	✓
	Mulga Lands	Charleville	✓	✓	✓
		Cunnamulla	✓		✓
		Bollon		✓	✓
		Quilpie	✓		✓
		Hungerford		✓	✓
	Mitchell Grass Down	Blackall	✓	✓	✓
		Winton	✓		✓
		Longreach		✓	✓
		Julia Creek	✓		✓
		Hughenden		✓	✓
	Desert Uplands	Barcaldine	✓	✓	✓
	Channel Country	Windorah	✓	✓	✓
	Eastern	Not stratified	Inglewood		✓
Emerald				✓	✓
Charters Towers				✓	✓
Western	Not stratified	Duchess	✓		✓
		Cloncurry	✓		✓
		Cloncurry	✓		✓

Appendix 2 Densities per km² of the commercially harvested macropod species 2004–2017

Eastern grey kangaroo														
Block	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Barcaldine	20.61	13.17	22.77	17.65	23.15	29.50	12.871	23.92	24.09	44.10	24.96	19.31	14.71	13.65
Blackall	7.57	7.10	6.22	7.51	8.28	11.19	7.082	6.08	9.87	19.41	10.59	8.75	5.29	4.97
Bollon	25.66	25.31		30.53		31.74	30.143		47.2		32.01	24.95		27.58
Charleville	17.51	19.91	15.96	12.05	11.20	12.95	12.229	28.11	25.12	26.77	11.77	8.4	10.47	9.32
Charters		1.63		5.02		5.33	5.568		3.37		3.14	2.01		1.53
Cloncurry	0.01		0.16		0.02		0.214	0.012		0.07		0.00	0.06	
Cunnamulla	13.20		9.97		11.44		11.642	32.82		41.04		35.87	18.73	
Duchess	0.00		0.00		0.00		0.00	0.00		0.00		0.00	0.00	
Emerald		3.95		3.41		4.05	5.035		2.75		7.01	5.29		7.88
Hughenden	0.77	0.58		1.16		0.97	0.795		0.53		1.17	1.41		1.01
Hungerford	1.16	1.10		0.77		0.94	0.651		2.20		4.00	3.79		3.36
Inglewood		8.72		18.62		9.75	12.326		29.10		32.73	49.88		42.02
Injune	13.53	18.85		18.62	6.82		16.599	17.75		31.32		22.52	40.62	
Julia Creek	1.08	0.87	1.05		0.76		0.276	0.28		0.84		0.84	0.2	
Longreach	9.05	8.48		6.63		6.61	6.129		18.07		20.17	5.25		3.85
Quilpie	1.86		0.97		1.42		2.795	1.57		3.61		4.65	2.66	
Roma	25.05	24.98	25.46	25.12		23.43	19.298		27.16		40.56	32.23		31.74
Taroom	8.12	13.37		8.44		7.87	7.362		14.98		13.24	12.65		8.19
Westmar	25.53		23.17		21.18		22.083	37.25		62.54		77.9	66.07	
Windorah	1.58	2.69	1.14	1.39	2.39	1.26	0.858	2.68	1.24	1.80	0.79	1.02	2.13	1.34
Winton	4.86	2.98	3.74		4.78		2.432	3.57		6.61		5.79	4.46	

2018 Quota Submission for Commercially Harvested Macropods in Queensland

Red kangaroo														
Block	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Barcaldine	6.07	4.07	11.07	6.72	9.03	9.83	7.58	10.05	7.83	12.30	9.2	16.24	8.54	8.25
Blackall	3.99	3.29	4.55	3.78	6.45	7.24	4.70	12.37	14.17	17.47	10.35	11.58	9.69	8.52
Bollon	4.13	8.87		8.35		11.16	9.90		7.78		7.27	9.83		8.87
Charleville	4.55	5.48	7.36	9.57	7.58	8.47	6.46	14.69	5.53	7.03	4.97	5.06	6.32	3.75
Charters		0.02		0.05		0.00	0.70		0.24		0.21	0.62		0.59
Cloncurry	2.14		4.18		6.17		3.01	3.34		5.91		4.06	3.45	
Cunnamulla	3.54		4.59		9.02		10.65	18.27		28.76		27.29	16.54	
Duchess	2.92		0.87		1.78		0.85	0.71		0.82		1.66	1.61	
Emerald		0.00		0.00		0.00	0.02		0.05		0.00	0.00		0.00
Hughenden	1.97	1.59		1.59		1.29	0.92		2.22		2.67	1.52		2.62
Hungerford	1.04	2.57		3.90		4.41	2.60		7.01		8.75	9.7		9.83
Inglewood		0.00		0.00		0.00	0.50		0.00		0.00	0.00		0.00
Injune	1.02	0.14	0.05		0.72		0.00	0.86		0.04		0.00	0.11	
Julia Creek	4.08	5.13	4.91		5.39		3.16	3.30		8.10		5.6	4.58	
Longreach	9.53	11.86		11.33		14.71	12.24		14.43		19.26	4.79		15.90
Quilpie	2.19		1.39		5.13		2.06	4.70		9.80		9.51	12.27	
Roma	2.19	1.62	2.54	2.66		2.37	2.26		3.47		5.02	2.83		2.96
Taroom	0.02	0.37		0.00		0.00	0.00		0.00		0.00	0.00		0.00
Westmar	0.97		0.59		0.55		1.14	1.49		1.14		2.28	4.69	
Windorah	4.42	4.52	7.32	4.48	9.85	12.62	6.67	10.47	10.77	11.84	8.11	12.29	12.16	16.95
Winton	3.69	5.02	5.62		6.05		3.32	4.44		9.73		16.98	8.57	

2018 Quota Submission for Commercially Harvested Macropods in Queensland

Common wallaroo														
Block	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Barcaldine	6.59	12.30	8.63	16.42	19.98	16.20	5.521	8.306	5.53	9.02	4.69	6.79	4.89	3.57
Blackall	18.02	21.17	22.15	34.98	39.14	49.05	23.819	21.549	20.2	54.43	28.58	24.86	11.29	9.32
Bollon	4.72	2.64		1.55		1.67	0.363		2.44		0.7	0.31		2.38
Charleville	7.21	6.20	5.79	4.66	4.70	5.36	11.803	13.439	10.65	5.09	4.47	4.07	3.81	1.24
Charters		0.30		2.84		0.61	1.069		0.51		1.03	0.22		0.23
Cloncurry	0.00		0.30		0.64		0.507	0.260		0.14		0.02	0.21	
Cunnamulla	1.68		0.45		0.64		1.951	0.611		0.70		2.53	2.60	
Duchess	0.57		0.11		1.69		0.743	0.093		1.16		0.00	0.00	
Emerald		0.02		0.00		0.78	0.024		0.33		0.19	0.32		0.25
Hughenden	1.65	1.28		2.28		0.24	0.411		0.94		0.93	0.22		1.59
Hungerford	1.19	0.36		0.24		0.48	0.254		0.47		1.27	1.93		2.33
Inglewood		3.08		4.03		0.34	1.013		1.22		3.18	3.42		11.16
Injune	0.01	2.30	1.34		0.91		4.053	0.00		0.63		0.84	0.60	
Julia Creek	2.74	0.00	0.04		0.11		0.013	0.00		0.00		0.00	0.03	
Longreach	17.96	21.57		18.59		12.69	9.185		17.77		15.67	8.84		4.17
Quilpie	5.41		0.78		3.36		2.686	3.007		5.58		7.51	3.42	
Roma	1.35	3.74	2.49	2.08		1.16	3.447		0.87		1.01	0.75		1.45
Taroom	0.22	2.04		0.17		1.05	0.253		0.02		0.38	0.18		0.35
Westmar	0.74		0.02		0.13		0.299	0.00		0.01		0.00	0.28	
Windorah	2.14	2.30	1.81	2.72	3.03	3.07	2.422	3.185	3.32	5.29	2.82	1.46	4.86	4.12
Winton	1.73	1.78	1.70		3.14		0.955	4.191		6.35		0.76	1.14	