

Statewide Landcover and Trees Study

Methodology overview v1.1



Queensland
Government

Prepared by: Remote Sensing Sciences, Department of Environment and Science

© State of Queensland, 2022.

The Department of Environment and Science acknowledges Aboriginal peoples and Torres Strait Islander peoples as the Traditional Owners and custodians of the land. We recognise their connection to land, sea and community, and pay our respects to Elders past, present and emerging.

The department is committed to respecting, protecting and promoting human rights, and our obligations under the Human Rights Act 2019.

The Queensland Government supports and encourages the dissemination and exchange of its information. This work is licensed under a Creative Commons Attribution 4.0 International License.



Under this licence you are free, without having to seek our permission, to use this publication in accordance with the licence terms. You must keep intact the copyright notice and attribute the State of Queensland as the source of the publication.

For more information on this licence, visit <https://creativecommons.org/licenses/by/4.0/>

Disclaimer

This document has been prepared with care, based on the best available information at the time of publication. The department holds no responsibility for any errors or omissions within this document. Any decisions made by other parties based on this document are solely the responsibility of those parties

If you need to access this document in a language other than English, please call the Translating and Interpreting Service (TIS National) on 131 450 and ask them to telephone Library Services on +61 7 3170 5470.

This publication can be made available in an alternative format (e.g. large print or audiotape) on request for people with vision impairment; phone +61 7 3170 5470 or email <library@des.qld.gov.au>.

Citation

Queensland Department of Environment and Science. 2022. Statewide Landcover and Trees Study – Methodology Overview v1.1. Queensland Government, Brisbane.

November 2022

List of acronyms

CNN	Convolutional Neural Network
CRF	Conditional Random Fields
DES	Department of Environment and Science
DMC-3	Disaster Monitoring Constellation-3
EDS	Early (Clearing) Detection System
ESA	European Space Agency
ETM+	Enhanced Thematic Mapper Plus
FPC	Foliage Projective Cover
GBR	Great Barrier Reef
JRSRP	Joint Remote Sensing Research Program
MCCV	Monte Carlo Cross Validation
NDVI	Normalised Difference Vegetation Index
NSW	New South Wales
NVIS	National Vegetation Information System
OLI	Operational Land Imager
QLUMP	Queensland Land Use Mapping Program
RSS	Remote Sensing Sciences (formerly Remote Sensing Centre in DES)
SLATS	Statewide Landcover and Trees Study
TM	Thematic Mapper
USGS	United States Geological Survey
VMA	<i>Vegetation Management Act 1999</i>

Glossary

Albers equal-area projection

A conic map projection which preserves the area of features. This projection is suitable for use when calculating the area of features.

Clearing activity

Human activity which results in the full or partial removal or destruction of woody vegetation from an area.

Foliage projective cover (FPC)

FPC is defined as the fraction of ground covered by the vertical projection of photosynthetic foliage of all strata (Specht, 1983). FPC is a metric that is used in remote sensing (i.e. satellite-based monitoring) as a direct estimate of the foliage (or leaves) on vegetation when viewed (vertically or near-vertically) from above, as is the perspective of the satellite. Herein, FPC refers to the foliage of woody plants only and is expressed as a percentage where: 0% FPC implies there is no woody plant foliage cover; and 100% FPC implies total or complete woody plant foliage cover.

Full clearing

A human-induced clearing event that results in the complete removal or destruction of woody vegetation, converting an area from woody to non-woody (i.e. less than 10% woody crown cover remains). These are areas that were mapped as woody in the woody extent map, but the clearing activity has sufficiently removed or destroyed enough woody vegetation to render the location non-woody, thus removing them from the woody extent map. These areas are included as one of the categories of clearing activity in SLATS reporting.

Geometric correction

Also referred to as geo-referencing, this process is used to accurately register satellite images to a ground coordinate system.

Image composite

An image composite refers to the multi-temporal compositing of image scenes. In the SLATS process, image composites are primarily used to address incomplete coverage of a Sentinel-2 tile on any one date due to the satellite orbital path, or to replace cloudy areas from one date with clear data from another date to maximise the useable data per scene.

Image mosaic

An image mosaic, as referred to in the SLATS process, is produced by combining multiple individual image scenes to produce a single seamless mosaic for the state of Queensland.

Partial clearing (major)

A human-induced clearing event that results in the partial but significant removal or destruction of woody vegetation. These are areas where greater than 50% of the woody vegetation has been affected by clearing but the area remains woody (i.e. greater than 10% crown cover remains). These areas are included as one of the categories of clearing activity in SLATS reporting but remain in the woody extent.

Partial clearing (minor)

A human-induced clearing event that results in partial, minor removal or destruction of woody vegetation. These are areas where the woody vegetation has been modified but less than 50% of the area has been affected by clearing, and it remains woody (i.e. greater than 10% crown cover remains). These areas are included as one of the categories of clearing activity in SLATS reporting but remain in the woody extent.

Radiometric standardisation

Refers to the process of correcting satellite imagery for atmospheric effects, seasonal differences in reflectance due to sun-sensor-ground geometry, and sensor characteristics. This standardisation or correction is particularly important for image mosaicking and comparing images over multiple time periods.

Woody plants

A plant that produces wood as its primary structural tissue. Woody plants may be trees, shrubs or lianas and are usually perennial.

Woody baseline

The 2018 map of woody vegetation extent (greater than 10% crown cover and minimum patch size of 0.5ha), which

forms the basis for SLATS woody vegetation monitoring, accounting and reporting.

Woody vegetation

Assemblages of woody plants. This includes stands of native vegetation, regrowth following clearing, plantations of native and exotic species, and woody weeds.

Contents

List of acronyms.....	iii
Glossary.....	iv
1 Introduction.....	1
1.1 Background.....	1
1.2 Objectives of SLATS.....	1
1.3 Recent changes to SLATS methodology.....	1
1.4 Purpose of this document.....	1
1.5 Independent peer review.....	1
2 The SLATS monitoring and reporting framework.....	2
2.1 Scope and key definitions.....	2
2.1.1 In scope.....	2
2.1.2 Out of scope.....	2
2.1.3 Key definitions.....	3
2.2 Overview of the SLATS monitoring, accounting, and reporting framework.....	4
3 Methods.....	6
3.1 Systems and data.....	6
3.1.1 Systems.....	6
3.1.2 Satellite imagery and imagery pre-processing.....	6
3.1.3 Field data.....	7
3.2 Woody extent mapping.....	8
3.2.1 Overview.....	8
3.2.2 Mapping specifications.....	8
3.2.3 Data.....	8
3.2.4 Automated classification.....	8
3.2.5 Manual editing and refinement.....	9
3.2.6 Data products.....	11
3.3 Woody vegetation clearing mapping.....	11
3.3.1 Overview.....	11
3.3.2 Mapping specifications.....	11
3.3.3 Data.....	11
3.3.4 Automated clearing detection.....	12
3.3.5 Clearing mapping.....	13
3.3.6 Data products.....	17
3.4 Woody vegetation age since last disturbance estimation.....	17
3.4.1 Overview.....	17
3.4.2 Data.....	17
3.4.3 Methods.....	18
3.4.4 Limitations.....	19
3.4.5 Data products and outputs.....	19
3.5 Woody vegetation regrowth mapping.....	20

3.5.1	Overview	20
3.5.2	Methods	20
3.5.3	Data products and outputs.....	22
3.6	Woody vegetation density estimation using foliage projective cover	22
3.6.1	Overview	22
3.6.2	Data.....	22
3.6.3	Methods	23
3.6.4	Data products and outputs.....	23
4	Data integration, reporting and accounting	24
4.1	Data integration.....	24
4.1.1	Annual updating of the woody extent.....	24
4.2	Reporting	25
4.2.1	The reporting package	25
5	References	27
6	Appendix.....	29
6.1	Landsat continuity: a sample-based estimation of woody clearing in 2018–19 using SLATS Landsat-based methodology.....	29
6.1.1	Introduction	29
6.1.2	Methods	29
6.1.3	Results	31
6.1.4	Conclusion	32

1 Introduction

1.1 Background

The Statewide Landcover and Trees Study (SLATS) is a scientific monitoring program undertaken by the Department of Environment and Science in partnership with the Joint Remote Sensing Research Program (JRSRP) and the Queensland Herbarium. The program works closely with the Department of Resources which administers the *Vegetation Management Act 1999* (VMA).

SLATS uses satellite imagery and field data to monitor and report changes in woody vegetation extent in Queensland and provide information about other woody vegetation attributes such as foliage density and age.

Data, reporting and other information from SLATS supports the VMA and a range of environment, natural resource and disaster management policy requirements and applications. This includes protection and management of the Great Barrier Reef (GBR), State of the Environment reporting, the Regional Ecosystem mapping framework, biodiversity conservation, fire management and planning, and natural capital and environmental accounting initiatives.

1.2 Objectives of SLATS

The primary objective of SLATS is to monitor, map and account for changes in the extent of woody vegetation across the state of Queensland on an annual basis. This includes documenting (i.e. baselining) the current extent and monitoring and mapping changes to that extent due to human-induced land clearing and natural or human-induced regrowth.

A secondary objective of SLATS is to provide additional data and information about the type of clearing activity and its purpose, and about the density and age of the vegetation that currently exists, is being cleared, or is regrowing. Where possible, SLATS will also map losses or disturbances to woody vegetation through natural causes such as cyclone damage or fire.

SLATS is complemented by other land cover change monitoring for ground cover and fire scars, and the Early Detection System—a regular, targeted, monitoring and proactive compliance tool which also supports the VMA.

1.3 Recent changes to SLATS methodology

SLATS monitoring and reporting methodology, up to and including the 2017-18 report, was primarily based on Landsat satellite imagery and focussed on woody vegetation clearing mapping. From 2018-19 onwards, SLATS has used a completely revised methodology that is primarily based on Sentinel-2 satellite imagery. This methodology incorporates monitoring components in addition to clearing mapping which includes maintaining a woody vegetation baseline, and monitoring and mapping new woody vegetation regrowth. The methodology for the revised program is the subject of this document.

Due to the revision of the SLATS methodology, including the transition from Landsat to Sentinel-2 satellite imagery, SLATS reporting from 2018-19 will not be directly comparable with previous reporting. However, as part of SLATS revisions a separate scientific study was undertaken to provide clearing estimates that were more comparable with previous SLATS reporting. An overview of the methods developed for that study are provided in the Appendices (Section 6).

1.4 Purpose of this document

The purpose of this document is to provide an overview of the current SLATS methodology which is based primarily on Sentinel-2 satellite imagery.

1.5 Independent peer review

The current SLATS methodology was the subject of a comprehensive independent peer review in 2021. The review was led by CSIRO and included experts in remote sensing, ecology, and natural resource management.

The review considered the science underpinning the program and its effectiveness in supporting environment and natural resource information requirements. It found that the methods are appropriate and consistent with best practice scientific reporting, and that the program will have ongoing impact for environment and natural resource management in Queensland.

2 The SLATS monitoring and reporting framework

2.1 Scope and key definitions

2.1.1 In scope

In general, SLATS monitors woody vegetation in line with the definition of vegetation in the VMA, which defines vegetation as woody native trees or plants, except for mangroves which are protected under other legislation. Due to the difficulty of discriminating native and non-native woody vegetation in the landscape using satellite imagery, non-native woody vegetation change is also included in SLATS monitoring and reporting (refer to Section 2.1.2.1 for details regarding this).

The present scope of SLATS is to map the current extent of woody vegetation in Queensland and monitor and map changes to that extent due to human-induced land clearing and natural or human-induced regrowth. This is limited by what can be reliably identified and mapped in Sentinel-2 satellite imagery, but also informed by other data sources, including Landsat time-series and very high-resolution satellite imagery.

SLATS also aims to determine the type of clearing activity and its purpose and estimates of the density of the vegetation that currently exists, is being cleared, or is regrowing. Time-series methods are also used to estimate the time since the woody vegetation was last disturbed and, where possible, to detect the time since it had started to regrow following a disturbance event.

The initial baseline woody extent aimed to capture all woody vegetation as at (nominally) August 2018 that has a crown cover of 10% and above (approximately 5-6% FPC) and a minimum patch size of 0.5ha. This includes woody vegetation in very sparse, sparse, mid-dense and dense classes (Table 1). Thereafter, all monitoring and mapping of changes to the woody extent (i.e. due to clearing and regrowth) are mapped for the same crown cover criteria but using a minimum patch size of 0.25ha.

Table 1: Overview of structural (vegetation density) formations for woody vegetation as used in SLATS reporting. Adapted from Scarth *et al.*, 2019.

Foliage Projective Cover (FPC) or Crown Cover (CC)			
Very Sparse/Isolated <10% FPC 0.25-20% CC	Sparse 10-30% FPC 20-50% CC	Mid-dense 30-70% FPC 50-80% CC	Dense 70-100% FPC >80% CC
tall open woodland	tall woodland	tall open forest	tall closed forest
open woodland	woodland	open forest	closed forest
low open woodland	low woodland	low open forest	low closed forest
tall open shrubland	tall shrubland	open scrub	closed scrub
low open shrubland	low shrubland	open heath	closed heath

2.1.2 Out of scope

The present scope of SLATS excludes some land cover, land cover change, and ecological attributes from the monitoring and reporting. This is due to a range of factors including the objectives for SLATS, the limitations of current scientific research to identify and accurately map some attributes state-wide, or because other programs in government are already mapping or documenting the attributes. SLATS may address some of these where appropriate in the future, and the program also maintains close ties with other state and commonwealth government programs to minimise duplication and ensure complementarity and information sharing.

Current exclusions from SLATS are briefly outlined in the following subsections.

2.1.2.1 Distinguishing native woody vegetation from non-native woody vegetation

SLATS mostly monitors native woody vegetation change but due to the difficulty of separating native and non-native woody vegetation using satellite imagery, especially woody weed species such as Prickly acacia (*Valchellia nilotica*), Parkinsonia (*Parkinsonia aculeata*) and Lantana (*Lantana camara*), reporting for some regions or localised areas will likely include some non-native woody vegetation change.

2.1.2.2 Non-woody vegetation change monitoring

The extent, density or change in non-woody vegetation communities is not currently monitored by SLATS. Refer to the [Queensland Ground Cover Monitoring Program](#) for further information regarding how non-woody vegetation is monitored by this program.

2.1.2.3 Vegetation height

Vegetation height is not currently considered, as it is not currently possible to reliably estimate height from optical satellite imagery such as Sentinel-2 or Landsat. This may be improved in the future through research into the use of radar, LiDAR and combinations of these sensors with optical imagery.

2.1.2.4 Vegetation composition

This information is available in the Queensland Herbarium's Regional Ecosystem mapping for Queensland.

2.1.2.5 Vegetation densification (i.e. thickening)

Vegetation densification, sometimes also referred to as thickening, is not presently monitored. This is an area of ongoing research in RSS/JRSRP. The current FPC product is intended to provide contextual attribution of the vegetation density to assist understanding of the types of ecosystems which are being cleared or regrown. While larger increases in the FPC product over time for a particular location may reflect real densification processes on the ground, insufficient research and validation of this product has been undertaken to reliably use it for tracking change in density, as these changes are often quite subtle and within the confidence intervals (i.e. error) of the FPC product. It is worth noting that the process of encroachment—the colonisation of a previously non-woody area by woody vegetation—will be monitored as a change in woody vegetation extent as part of the regrowth monitoring, where the change can be reliably identified and mapped following SLATS mapping specifications.

2.1.2.6 Fire

Fire-affected areas are presently mapped separate to SLATS. Refer to the [Queensland Fire Scar Mapping Program](#) for further information. Where SLATS can identify human-induced clearing associated with a fire event, this is mapped. However, fire-affected areas are generally assumed to be temporary, non-anthropogenic changes in woody vegetation, even if the ignition source was human induced. It is important to note that some estimates of vegetation age may be affected by historical fire events; regrowth age estimates may be influenced by fire and other natural disturbances which impact the canopy significantly, even if temporarily. If a fire or other natural disturbance such as cyclone damage, flood etc. causes the modelled woody probabilities to fall below a given threshold and this causes a break point, regrowth tracking is reset, and age would be determined from the 'next' period of regrowth/recovery detected after that point in time (depending on the other heuristic conditions—refer to Section 3.4.3 for further details).

2.1.2.7 Land use and land use change

Land use is mapped by the [Queensland Land Use Mapping Program](#). SLATS includes some information related to land use—the *landcover replacement class* (see Section 3.3.5.2)—but this is only intended to be indicative of the land use for which a clearing activity has been undertaken.

2.1.2.8 Other natural or non-human-induced change

SLATS attributes natural or non-human-induced changes as part of the clearing mapping process where they are possible to identify and map, but these changes are not included in any reporting. This includes change due to natural disaster impacts (e.g. cyclones, floods, droughts) and other natural tree death (e.g. senescence, dieback).

2.1.3 Key definitions

The following are key definitions as they apply in the context of the SLATS monitoring, reporting, and accounting framework. The definitions are therefore defined by what can be reliably identified and mapped consistently over large areas using (mostly) satellite imagery. For other terms and definitions used in this document, refer to the glossary on page iv.

2.1.3.1 Woody vegetation

Assemblages of woody plants with greater than 10% crown cover. These may be trees, shrubs or lianas, and are usually perennial. An assemblage may include uncleared native vegetation, regrowth following a previous clearing event (or events), plantations of native and exotic species, and woody weeds.

2.1.3.2 Woody vegetation clearing activity

The anthropogenic (i.e. human-induced) removal or destruction of woody vegetation. This may be 'full clearing' (i.e.

a conversion from woody to non-woody), or 'partial clearing' (i.e. there has been some clearing activity for a given location but it is not sufficient to render the area non-woody). Refer to Section 3.3 for details.

2.1.3.3 Woody vegetation age since disturbance

The estimated time since the woody vegetation was last significantly disturbed, or where possible to detect, the time since it started to grow or regrow following a disturbance. Estimates are based on time series modelling of woody vegetation and SLATS change history within the period 1988 to present. Refer to Section 3.4 for details.

2.1.3.4 Woody vegetation regrowth (or regrowth)

Woody vegetation that has regrown due to natural or human-induced processes and is determined to be sufficiently woody to be added to the woody extent data set (i.e. meets the criteria of having 10% crown cover and stand area of >0.25ha). Refer to Section 3.5 for details. As noted in Section 2.1.2.5, regrowth can include areas where encroachment has occurred.

2.1.3.5 Woody vegetation density

The estimated density of woody vegetation's foliage cover based on a data product that estimates FPC and is derived from Sentinel-2 satellite imagery, calibrated by field estimates of FPC (refer to Section 3.6 for details).

2.2 Overview of the SLATS monitoring, accounting, and reporting framework

Figure 1 is a schematic diagram of the key components and timeline which forms the basis of the SLATS monitoring, accounting, and reporting framework. The current framework is based on a conceptual model of establishing a detailed baseline account of the woody vegetation extent, age since disturbance and density for the state (as of 2018), and then monitoring and accounting changes to that extent due to clearing and regrowth, reporting annually. Estimates of woody vegetation age since disturbance and density are also updated annually to inform reporting. Section 3 provides details about the methods and workflows which support the framework.




Figure 1 Schematic example of the SLATS monitoring, accounting, and reporting framework

3 Methods

3.1 Systems and data

SLATS is supported by an extensive computing infrastructure and comprehensive archive of data. Data and programs (i.e. algorithms or code) used and produced are subject to quality control systems and standard operating procedures used by DES's RSS for image and field data processing, storage, and management. The systems and processes used are fully documented in a WIKI-based system and include peer-reviewed processes. All image processing follows internationally accepted and/or published standards. Data sourced from external parties are not incorporated into the workflow without appropriate metadata, including lineage statements.

3.1.1 Systems

All data processing is undertaken on the high-performance computing (HPC) facility at the Ecosciences Precinct, Brisbane. The HPC incorporates a multi-node Linux cluster, a mass storage on-line disk and near-line tape silo which houses the image archive and downstream products and is connected to the high-speed national science network portal (AARNet) enabling efficient downloading of large data sets.

Image metadata, field and other spatial data sets are stored in RSS's PostgreSQL spatial database, which is integrated with the image archive and processing systems to enable efficient querying of /access to SLATS data sets. Imagery and other data sets are governed by a naming convention which is systematic, structured, and descriptive, and supports automated processing (Flood and Danaher, 2013).

Processing is largely done using open-source software with much of the code written using a Python base and open-source or in-house developed libraries. All code written for SLATS is maintained in Git version control repositories, hosted by Gitlab.

3.1.2 Satellite imagery and imagery pre-processing

The primary satellite data used in the various components of the current SLATS program include imagery from: Landsat TM, ETM+ and OLI, Sentinel-2A and -2B, and DMC-3 TripleSat (Earth-i).

3.1.2.1 Sentinel-2 imagery

Sentinel-2 MSI imagery (10m/20m) from ESA's Copernicus program is used by SLATS as the primary image data for ongoing annual monitoring and reporting of woody vegetation clearing (Section 3.3), regrowth (Section 3.5), and density (Section 3.6). It is also used together with Earth-i and other very high-resolution data to inform woody extent refinement (Section 3.2).

Level 1C Sentinel-2 image products are routinely downloaded from the Australasian Regional Copernicus Data Hub (<http://www.copernicus.gov.au/>) and then processed to surface reflectance as described in Flood, (2013a). The imagery is geometrically corrected by ESA. Currently, ESA's geometric correction process is fully operational with geolocation accuracy reported to be within one 10m pixel. However, in the early part of the time series (2016-2019) ESA's quality tracking shows variable performance with long-term average location accuracy closer to 11m at 95% confidence (exceeding one 10m pixel; ESA, 2019). SLATS therefore undertook an additional geometric co-registration step of end-date image to start-date image for a small number of image pairs used in SLATS clearing detection. This was required to reduce misregistration effects in the clearing index. For processing of indices developed for Landsat such as FPC, fractional cover and the SLATS clearing index, the Sentinel-2 20m (Landsat-analogous) short-wave infrared (SWIR) bands are resampled to 10m using cubic convolution. Processing of these indices is described in the relevant sections. Cloud and cloud shadow masks are computed using the methods of Zhu *et al.*, (2015). Cloud and shadow masks are no longer used in the production of the SLATS clearing index as cloud-free image selection is almost always possible and current cloud mask performance is variable for Sentinel-2 imagery.

RSS has complete holdings of Sentinel-2A and -2B imagery acquired over Queensland from late 2015 and ongoing. From 2017 onwards this represents a five-day return interval.

3.1.2.2 Landsat imagery

The long time series of Landsat imagery (1988-ongoing) is used as the primary image data to derive temporal woody vegetation age since disturbance estimates (Section 3.4), It is also used extensively as an ancillary data set to provide historical context in SLATS change detection (Section 3.3) and woody extent refinement (Section 3.2),

and to augment the shorter Sentinel-2 time series in regrowth detection (Section 3.5).

RSS acquires Collection-2 Landsat-8 and -9 satellite imagery (combined represents an eight-day return interval over all areas of Queensland) from the USGS via a nightly automatic process that queries all new imagery using the USGS Machine-to-Machine (M2M) API tool (<https://m2m.cr.usgs.gov/>) and then downloads the data from their Amazon Web Services (AWS) object storage service (S3). Prior to this, Collection-1 Landsat data was downloaded directly from the USGS Earth Explorer site. RSS ceased downloads of Landsat-7 on 30 November 2021 as Landsat-9 data came fully online. Landsat-7 has the SLC-off issue and was considered redundant given availability of Landsat-8 and -9 in addition to Sentinel-2A and 2B data. Landsat imagery is geometrically corrected by the USGS. Analyses by the USGS suggest that the locational error is below a single pixel (Storey *et al.*, 2014). SLATS uses the USGS geometric correction, without modification. All Landsat data are processed to surface reflectance as described in Flood (2013a). Cloud and cloud shadow masks are computed using the *Fmask* methods of Zhu and Woodcock (2012).

RSS holds the complete archive of historic USGS Landsat imagery from Landsat-5 TM, -7 ETM+ (up to 30 November 2021), -8 OLI, and -9 OLI-2 acquired over Queensland during the period 1988 to present.

3.1.2.3 Seasonal fractional cover composites

SLATS uses time series of seasonal composites of fractional cover (Scarath, 2008) for woody vegetation age since disturbance estimates (Section 3.4) and regrowth detection (Section 3.5).

Three-monthly seasonal fractional cover composites are produced for Queensland for both Landsat (30m) and Sentinel-2 (10m) processing streams using the medoid method described in Flood (2013b). Prior to compositing, topographic shadow, cloud and cloud shadow masks are applied.

3.1.2.4 Earth-i DMC-3 Imagery

High-resolution (~80cm) satellite image mosaics of Queensland were purchased by the Queensland Government for the years 2016, 2017 and 2018. The 2017 and 2018 mosaics were the primary image data used in the development of the woody extent (Section 3.2).

The data were purchased as three-band visible (blue, green, red) imagery, pan-sharpened from a nominal resolution of 3.2m to 80cm pixel size. The imagery was captured as many small tiles over a range of months, extending between April and November in each year, by the Disaster Management Constellation (DMC) program's three DMC-3 TripleSat satellites. For each year, the imagery was pan-sharpened, colour-balanced and mosaicked by Earth-i, and finally supplied as 32 large mosaic tiles covering the whole of Queensland, with some overlap between tiles. Once on the RSS systems, the imagery was resampled using cubic convolution resampling into Albers Equal Area projection to a pixel size of 1m. The DMC satellites are pointable, allowing for view angles up to 30 degrees off-nadir to avoid cloud contamination. This allowed for largely cloud-free image mosaics, however some small cloud and poor image quality effects, due in part to larger view angles, are evident in all the mosaics.

3.1.2.5 Ancillary data

SLATS uses a range of ancillary data to inform the operational mapping components. These data include:

- high resolution satellite imagery and aerial photography available through online image services such as Planet (QSat), Google Earth, One Atlas Living Library, and the Queensland Government's *Queensland Globe*
- complementary remote sensing products, for example DES's Sentinel-2 fire scar data, and the Northern Australia Fire Information's fire hotspots and fire scar maps (<https://www.firenorth.org.au/nafi3/>)
- airborne lidar and terrestrial laser scanner data
- regional ecosystem mapping.

3.1.3 Field data

RSS hosts a national field database, which incorporates a comprehensive collection of vegetation measurements collected across Queensland and nationally over a 20+ year period, encompassing much of the time that Landsat and Sentinel-2 satellites have been operating. Field data such as the star transect collection of vegetation cover measurements (Muir, 2011) are used to calibrate and validate the remote sensing products that SLATS uses and produces. Field data including measurements, observations and extensive photos are stored in RSS's PostgreSQL spatial database.

3.2 Woody extent mapping

3.2.1 Overview

A woody extent data set is required to inform ongoing operational mapping components (clearing and regrowth) and to form the baseline and basis for the woody vegetation change accounting framework of SLATS. The baseline woody extent data set and subsequent annual updates also serve as useful stand-alone data products for a range of other applications where detailed woody extent mapping is required, addressing a data gap for the government.

To develop the woody extent baseline, a data set showing the presence/absence of woody vegetation was derived from a Convolutional Neural Network (CNN) classification and the 2017 high resolution (re-sampled) Earth-i (1m) satellite imagery (Flood *et al.*, 2019). This initial output then underwent extensive manual refinement by SLATS scientists to produce a detailed map of woody vegetation extent for Queensland. The 2017 woody extent was subsequently manually revised and updated, informed by SLATS clearing data for the 2017-2018 monitoring period, and once available, the 2018 Earth-i image mosaic (refer Section 4.1.1.1). The resulting 2018 woody vegetation map established the baseline for SLATS ongoing monitoring, accounting, and reporting.

At the end of each annual SLATS monitoring period the woody extent is updated using the SLATS clearing and regrowth mapping for that period (refer Section 4.1.1).

3.2.2 Mapping specifications

The scale of the woody extent data set is intended to capture features visible at a nominal map scale of 1:10,000: stands of woody vegetation greater than 0.5ha with a crown cover greater than 10% are represented. A minimum width of 20m was applied to linear features. The 10% crown cover was chosen to ensure sparser vegetation in the rangelands of Queensland was represented, and to enable SLATS scientists to define extent boundaries consistently and accurately; delineating woody vegetation extent boundaries below this threshold can be ambiguous. Recognising that it is very challenging to estimate height in optical imagery, there is currently no height specification.

3.2.3 Data

The 2017 1m resolution Earth-I DMC-3 mosaic (Section 3.1.2.4) was used as the primary data source in the CNN woody classification and subsequent interpretation and manual refinement of the woody extent map.

It has a spatial resolution that allows a higher level of detail than is routinely feasible with moderate resolution satellite imagery from Sentinel-2 and Landsat. This is particularly important for mapping regions of sparse, open vegetation.

3.2.4 Automated classification

Convolutional Neural Networks (CNN) are a subset of machine learning algorithms that exploit patterns, texture, and shapes in images, and have been shown to outperform traditional classifiers. In particular, the U-net CNN (Ronneberger, 2015) was chosen as it is less reliant on extensive training data than other CNN approaches and results in a per-pixel classification of the original data. The U-net is structured around stepwise filtering and degrading of imagery to different resolutions and then upscaling back to original resolution, allowing detection of structures and textures at different scales.

3.2.4.1 Training data

The U-net model needs to be trained on many small patches of the source data, in this case the Earth-i imagery, each with a corresponding label image in which all pixels are labelled with the correct class: in this case, woody or non-woody. A patch size of 128 x 128 pixels was used. The patch size must be larger than the objects and textures to be detected. In this case, it needed to allow sampling whole groups of trees in sometimes sparse canopies, rather than single tree crowns.

As there was not an appropriate high-resolution woody/non-woody data set available, the required labels were derived from a set of 1km x 1km Earth-i image subsets, regularly sampled from a grid across each of the 32 mosaic tiles. After eliminating subsets with cloud or poor image quality and augmenting with targeted sampling in regions of under-represented, highly diverse land covers, a set of 827 image subsets were used to develop the set of woody/non-woody labels. A simple threshold-based classification of the Earth-i green band coupled with a binary closing filter was used to produce an initial woody/non-woody classification for each image subset. These were then manually refined to produce the set of correct woody/non-woody labels. The required training patches of 128 x 128 pixel were then extracted from each image-label pair in an overlapping grid and used in the model.

In the development of the woody extent map training data, an operator interpretation of woody vegetation as captured in high resolution imagery was required. It is worth noting that in high resolution imagery, a single tree crown can be captured by many pixels, conceptually different from representations of woody vegetation in moderate resolution satellite images from Sentinel-2 and Landsat, where a single pixel can be larger than a single tree.

No attempt was made to map individual tree crowns, but rather contiguous regions of woody vegetation cover which include tree/large shrub crowns, shadows, and the gaps between crowns, at a wide range of densities, broadly consistent with existing coarser scale mapping of regional ecosystems (Flood, *et al.*, 2019). These textures are the primary visual indicators of woody cover, at a range of densities. Biophysical quantities such as height were implicit in the human operator's view of the imagery, but are not always easily quantifiable from optical imagery, even with a pixel size of 1m.

3.2.4.2 Model performance

The model's ability to predict woody/non-woody pixels was assessed as approximately 90% accurate using Monte Carlo Cross Validation (MCCV), and with consistent performance across most of Queensland (no spatial bias). The MCCV involved 50 repetitions of the model fit, where in each iteration 70% of the training data subsets were used to fit the model and 30% withheld for validation. This means that in any model run, subsets which were held out were independent of the model. This gives a robust measure of accuracy from a number of accuracy assessments, each one conducted on an independent data set.

3.2.4.3 Woody classification

All training data were used to produce the final model, and the model then used to produce a 2017 woody classification for Queensland from the 1m resolution Earth-i mosaic tiles.

3.2.4.4 Downscaling for manual refinement

The 1m resolution classification was downscaled to 10m to better match the intended mapping scale, suitable for integration with Sentinel-2-based monitoring and to facilitate practical manual editing. In the downscaling, a 10m pixel was labelled as woody if $\geq 10\%$ of intersecting 1m pixels were labelled woody. A 10m binary closing filter was applied to connect small fragments that form part of a larger area/ecosystem (or stand) of classified woody areas, reduce the detail to manageable levels for editing, and to counter the errors associated with inconsistent mapping of individual crowns, and the effects of crown shading and illumination in the 1m data. As a final step, small clumps of woody vegetation and gaps (non-woody clumps) were filtered to implement the minimum mapping unit of 0.5ha.

3.2.5 Manual editing and refinement

3.2.5.1 Decision rules

Features were categorised as either woody or non-woody based on a set of decision rules informed by visual inspection of the data, the needs of existing data sets/programs, and expert knowledge. These decisions were based on considerations for SLATS and Spatial BioCondition and other vegetation monitoring applications, as well as ecological characteristics of the vegetation. No distinction was made between native and non-native vegetation—woody vegetation such as woody weeds and horticultural crops are included as woody in the final classification.

3.2.5.2 Manual editing

The data set was manually refined using a vector-editing approach in ArcGIS Pro software. A PostgreSQL distributed geodatabase was utilised to version data and enable reconciliation of edits to a master data set. Each scientist undertaking the editing was assigned a unique connection file to access the data set. Tiles comprised of a 25 x 25km systematic grid were used to sub-divide and manage the data set and to track mapping progress.

Each scientist was allocated an area to assess (subset by Queensland's bioregions) allowing for defined regions for editing so that specific experience and knowledge was developed of the features and vegetation types for the respective bioregion. Each scientist worked across a number of different bioregions over the course of the project, also building landscape knowledge to apply in other operational components of SLATS.

SLATS scientists visually checked mapping polygons against available satellite imagery (Earth-i 2017, Sentinel-2) at an on-screen scale of $\sim 1:10,000$. Misclassification or spatial (i.e. boundary) errors were edited using the suite of editing tools within the software package. These errors were generally caused by misclassification in the U-net modelling and could be due to factors attributable to the imagery (i.e. cloud, geometric/radiometric distortion, quality of imagery/colour balancing), or low accuracy in the modelling prediction. Where there was ambiguity in the classification of a feature, ancillary data was utilised to aid interpretation. Generally, ancillary data was limited to

regional ecosystem data (Queensland Herbarium 2019), land use data (QLUMP 2019), and previous SLATS clearing data. Expert opinion from senior botanists at the Queensland Herbarium was sought for specific areas of uncertainty or for specific vegetation types.

3.2.5.3 Mapping peer review and quality assurance (QA)

An in-house peer review process was established for the initial few months of the project to increase discussion between staff and achieve greater consistency between mappers in implementing decision rules. Scientists checked the editing of colleagues and provided feedback as comments in a point-based data set. Areas of disagreement or confusion were discussed across the wider team and decisions documented.

Additional to the initial peer review, a QA process was conducted by senior staff members throughout the entire editing phase to ensure accuracy and consistency. The QA process allowed senior staff to review all mapping with the goal of identifying and editing any obvious errors and providing feedback to editors in a timely manner. The process involved a systematic visual inspection of the edited outputs, similar to the original editing workflow but with a focus on final corrections and identification of woody vegetation, under- or over-represented, with regards to the mapping specifications.

3.2.5.4 Refinement through re-modelling

Certain features (i.e. urban, sparse vegetation, regrowth, mangroves) were not consistently predicted in the original CNN modelling resulting in excessive manual editing. Urban areas were particularly poorly classified and significantly over-estimated woody vegetation presence, likely due to shadowing effects and spectral/textural confusion from buildings and other infrastructure interspersed with woody vegetation and under-representation of urban areas in the training data set. Additionally, the generalised filtering processes used to degrade from 1m to 10m for manual refinement did not always adequately represent highly detailed land use/cover types as urban. For these areas, an iterative modelling approach was trialled using the U-net architecture (as described in Section 3.2.4) to create a more accurate baseline for further checking/editing. The trial was conducted using data from edited mapping in the Cairns urban region where the overestimated woody vegetation in urban areas had been removed by manual editing. This was then used to re-train a U-net model and predict woody vegetation in urban areas in Southeast Queensland. On visual inspection, outputs from the remodelled predictions in urban areas were more accurate and required significantly less manual editing. Based on this result, further urban regions across Queensland were re-modelled and used as the new baseline for further editing.

Re-modelling was also expanded to other areas where the original model predictions of woody vegetation were determined to be poor or inconsistent following visual inspection. These areas included: low woody regrowth in the Brigalow Belt; sparse patchy vegetation in the Mulga Lands; and mangroves and wetlands in Cape York and the Gulf Plains. Once available, the Earth-i 2018 imagery allowed for further remodelling for cloud-affected areas in the 2017 baseline. Training data for these re-modelling exercises was derived from the edited woody extent mapping within the same region as the area being re-modelled. Care was taken to ensure no vegetation change had occurred to ensure a consistency in the map currency. All re-modelled areas were then subject to the same editing and review process described in Sections 3.2.5.2 and 3.2.5.3.

3.2.5.5 Field work

An extensive field program was undertaken to provide SLATS scientists with the opportunity to calibrate their desktop image interpretation with on-ground observations. Largely qualitative observation, the field work facilitated a greater understanding of the vegetation types within and between bioregions improving consistency within the SLATS team in applying decision rules at the desktop. Field work also allowed for checking and refinement of the mapping. Areas of uncertainty or ambiguity in the mapping were accessed by vehicle where possible and observations of vegetation type, age, and condition were recorded, and field photographs were captured at the location if required. Further observations were captured opportunistically at points-of-interest *en route*. At the conclusion of each field trip, observations and photographs were compiled into a central database and used to further refine the mapping, where required.

3.2.5.6 Updating from 2017 to 2018 baseline

Upon completion of the 2017 woody extent map, it was subsequently updated to create the 2018 woody extent baseline using the SLATS clearing data for the 2017-18 monitoring period (detailed in Section 4.1.1.1) and filtered to maintain the 0.5ha minimum mapping specification (refer Section 4.1.1.2).

3.2.5.7 Successive annual updates

The woody extent is updated annually at the end of each SLATS monitoring period using the completed clearing (from 2018-19 onwards) and regrowth (from 2019-20 onwards) mapping for that period. The procedures for successive updating of the woody extent are detailed in Section 4.1.1.

3.2.6 Data products

The woody extent baseline and subsequent annual updates (refer to Section 4.1.1) are stored in one single vector data set, with attributes stored in a geodatabase that enable production of woody extent for a given year, or data about woody vegetation extent change between years, as well as a range of queries across multiple years and regions. SLATS has published the woody extent for 2018 (the baseline year) and 2019, the year which incorporates the first change monitoring period for this current methodology and will continue to publish woody extent updates in each successive annual monitoring period.

3.3 Woody vegetation clearing mapping

3.3.1 Overview

Historically, and ongoing, a state-wide clearing data set is required to inform policy evaluation for the VMA, as well as to service a range of land management and biodiversity conservation requirements across government. In terms of the current program's framework, it is also a fundamental component of the woody vegetation account, documenting the woody extent loss and adding to the non-woody areas which are monitored for regrowth in further monitoring periods. Woody vegetation clearing is defined as the removal or destruction of woody vegetation by human-induced (i.e. anthropogenic) means.

For over twenty years, and up until the 2017-18 monitoring period, SLATS has mapped the location and extent of woody vegetation clearing across the state of Queensland using Landsat Thematic Mapper (TM) and Enhanced Thematic Mapper (ETM+) imagery. The ability to detect woody vegetation change due to clearing, particularly in Queensland's sparser woodlands, is ultimately limited by the 30m spatial resolution of the Landsat imagery. SLATS has transitioned the Landsat approach to use Sentinel-2 satellite imagery at a spatial resolution of 10m to improve the detection and mapping of woody vegetation clearing in the current methodology.

The previous Landsat-based clearing mapping method detected woody vegetation change between two dry season Landsat images approximately twelve months apart. The mapping process involved two stages: (i) the automated generation of a woody vegetation clearing index that characterises reflectance differences between the two image dates and highlights possible clearing of woody vegetation as a probability measure (Scarth *et al.*, 2008); and (ii) extensive visual interpretation, manual refinement and quality assurance checking of the possible clearing detections to produce a final, high quality, clearing data set.

In general, the Landsat two-date method still applies to Sentinel-2 imagery, however there has been some necessary adaptation and modification of both the clearing index (detailed in Section 3.3.4) and the subsequent manual refinement approach (detailed in Section 3.3.5) necessitated by differences between Landsat and Sentinel-2 data (resolution, radiometrics and length of time series) and particularly the explicit incorporation of the woody vegetation extent data in the mapping process.

3.3.2 Mapping specifications

The Landsat woody vegetation clearing index has been adapted to apply to Sentinel-2 10m data and the manual refinement and quality checking is all undertaken at a scale suited to Sentinel-2's 10m spatial resolution.

The clearing mapping is undertaken (on-screen) at a scale of approximately 1:20,000. The minimum clearing size delineated is 0.25ha. As noted in Section 2.1.1 the 2018 baseline woody extent aimed to capture all woody vegetation as at (nominally) August 2018 that has a crown cover of 10% and above (approximately 5-6% FPC) and a minimum patch size of 0.5ha. This includes woody vegetation in very sparse, sparse, mid-dense and dense classes (Table 1). Thereafter, all monitoring and mapping of changes to the woody extent (i.e. due to clearing and regrowth) are mapped for the same crown cover criteria but using a smaller minimum patch size of 0.25ha.

These specifications are a compromise between requirements for the maintenance and updating of the woody extent data, deriving the benefits of the higher spatial resolution of Sentinel-2 data, and providing clearing data at a suitable level of detail to address a range of user requirements while maintaining consistency and efficiency for the annual program.

3.3.3 Data

3.3.3.1 Sentinel-2 imagery

The current clearing detection methods use Sentinel-2 imagery corrected to surface reflectance (Section 3.1.2.1). There are 245 Sentinel-2 tiles covering the state of Queensland. As in the Landsat monitoring approach, for each tile, a pair of Sentinel-2 surface reflectance images is used to represent the start and end dates of the 12-month monitoring period. The Sentinel-2 end date for the previous monitoring period becomes the start date for the current period. At the start of each monitoring period, the set of end date images covering Queensland are

manually selected to be as close as possible to a nominal dry season date of 15 August giving preference to minimal cloud cover, smoke, haze and greenness. Dry season imagery (July-Sept) tends to show the greatest contrast between woody vegetation and grass. Reasons for nominating the 15 August date are outlined in Section 4.2.1.5.

Approximately 60 of the 245 Sentinel-2 tiles are a composite of two dates (usually only a couple of days apart) due to the satellite flight path schedule. All SLATS dates are stored in a database for use by in-house scripts and general SQL queries.

3.3.3.2 Woody vegetation clearing index training data

The original Landsat woody vegetation clearing index was developed using training data derived from historic finalised SLATS clearing data mapped over numerous monitoring periods covering 2000-01 to 2010-11 (Scarath, Gillingham, and Muir, 2008). The original Landsat model was in part reliant on a long time series component, not applicable to the short Sentinel-2 time series. Nor was a large set of finalised Sentinel-2 clearing data available to train a new model directly on Sentinel-2 imagery.

Leveraging from work undertaken by NSW SLATS to adapt the original Landsat clearing index to SPOT imagery (Flood *et al.* 2020), a new form of the model (Equation 1c in Flood *et al.* 2020) was fitted on the original Landsat historic clearing training data set. The resulting index was then adapted to Sentinel-2 imagery using the radiometric relationships described in Flood (2017) to transform Sentinel-2 reflectance values to Landsat equivalents. This is possible given that the spectral bands are similar to those available from the Landsat sensor (Table 2).

Table 2: Resolution and spectral window for each of the corresponding bands for each sensor

Generic band name	Sentinel-2 MSI			Landsat 8 OLI		
	Pixel resolution (m)	Band number	Spectral wavelength (µm)	Pixel resolution (m)	Band number	Spectral wavelength (µm)
Green	10	3	543-578	30	2	530-590
Red	10	4	650-680	30	3	640-670
Near Infra-Red (NIR)	10	8	785-900	30	4	850-880
Short Wave Infra-Red 1 (SWIR 1)	20	11	1565-1655	30	5	1570-1650
Short Wave Infra-Red 2 (SWIR 2)	20	12	2100-2280	30	7	2110-2290

3.3.4 Automated clearing detection

The modelled woody vegetation clearing index generates values ranging between 0 and 1000 where 0 indicates no clearing and 1000 indicates a very high probability of clearing. In Scarath *et al.* (2008), Receiver-Operator Curve analysis of the Landsat clearing index was used to select six distinct thresholds resulting in a clearing classification based broadly on the likelihood of clearing whilst balancing omission and commission errors. A similar set of thresholds was applied to the Sentinel-2 based woody vegetation clearing index resulting in a set of six classes approximating varying likelihood of clearing from low through to very high. These classes are then used to inform mapping decisions by scientists in the manual refinement stage.

The woody extent for the start year in the monitoring period is used to identify any non-woody pixels in the clearing probability classes. For example, for 2018-19, this is the 2018 baseline woody extent. Ideally, pixels which are non-woody to start with would be eliminated from further scrutiny and the manual editing stage would only need to consider the woody areas for clearing assessment. Generally, the woody areas are still a focus of the mapping, however, at least in the early stages of the current methodology, an additional set of classes is used to identify potential clearing outside of the woody extent to ensure areas missed by the woody extent are still mapped if clearing has occurred. The woody and non-woody probability clearing classes are labelled with a set of codes listed

in Table 3.

The woody vegetation clearing index and change classification rasters are automatically generated for each of the 245 image pairs of start and end dates covering Queensland for the monitoring period.

Table 3: Clearing index thresholds in the change classification layer for woody and non-woody areas delineating classes ranging from low to increasing probability of clearing

Clearing index threshold	>420	370-420	320-370	270-320	220-270	170-220
Woody change classification codes	39	38	37	36	35	34
Non woody change classification codes	69	68	67	66	65	64

3.3.5 Clearing mapping

3.3.5.1 Mapping clearing using Sentinel-2 vs Landsat

Transitioning the woody vegetation clearing monitoring from medium resolution Landsat imagery with 30m pixels to high resolution Sentinel-2 imagery with 10m pixels involved a significant change in the scale of the monitoring. This in turn necessitated a change in the concept of clearing mapping. The medium resolution Landsat data was able to be used as a woody classifier (either modelled via, for example Foliage Projective Cover models, or by manual interpretation) since a single Landsat pixel tended to be larger than an individual shrub or tree crown. Each pixel captures both the crown and inter-crown gap of an area of woody vegetation. As resolution increases, this model starts to break down, since each pixel becomes a progressively purer representation of crown or non-crown (Zhang et. al. 2019). At higher and higher resolutions, a woody area then consists of both woody and non-woody pixels, particularly in ecosystems which are naturally more open. Given the scale change, an unbiased estimate of the area cleared changes from counting woody change pixels, to delineating the woody area (crown and inter-crown gap pixels) impacted by the clearing event. This area is referenced throughout this document as a “clearing footprint”. This is more representative of the total area impacted by the clearing event and more independent of the resolution of the sensor.

3.3.5.2 Attribution including ‘landcover replacement classes’

During the manual editing stage, each area of woody vegetation clearing is attributed with a *landcover replacement class* (Table 4). This provides an indication only of the land cover/land use purpose for which the vegetation was cleared or modified. Assignment of these classes is primarily based on visual interpretation, with reference to ancillary data sources. In areas where there are many different forms of land use, it can be difficult to interpret the final landcover replacement class and therefore this classification is indicative only. For example, land assigned to the class *pasture* may later be converted to *settlement* or other classes. Landcover replacement classes used in the Sentinel-2 change monitoring have been adapted from the Landsat-based clearing monitoring.

Table 4: SLATS replacement landcover

Replacement land cover	Description
Pasture	Grazing and other general agricultural land management practices—this class includes clearing for pasture, internal property tracks, fence lines or fire breaks.
Crop	Cropping or horticultural purposes.
Forestry	Timber harvesting in state or privately owned native or exotic (e.g. pine) forests or plantations.
Mine	Mining activities (including coal seam gas infrastructure).
Infrastructure	Utilities such as roads, railways, water storage, pipelines, powerlines etc.
Settlement	Urban development.

Additional attribution discriminates the type of clearing activity in terms of how much destruction or modification of the vegetation has occurred for any clearing event. This serves two purposes: (i) areas which are partially modified are able to be reported as such, thus providing a more nuanced component to the reporting of the clearing areas – this is particularly relevant for those regions where partial clearing is routinely undertaken under VMA codes or in unregulated areas for routine agricultural land management (e.g. fodder clearing and harvesting in the Mulga Lands); and (ii) it enables updating of the woody extent after each monitoring period such that partially cleared areas remain in the woody extent and are therefore considered in future monitoring.

Three classes are used to represent the degree of modification:

- i. Full clearing refers to clearing areas where the clearing activity has resulted in conversion from woody (i.e. greater than 10% crown cover) to non-woody (i.e. less than 10% crown cover). Full clearing will result in removal from the woody vegetation extent data set.
- ii. Partial clearing (major) refers to clearing areas where greater than 50% of the patch area has been removed and greater than 10% crown cover remains. These areas remain in the woody extent.
- iii. Partial clearing (minor) refers to where less than 50% of the clearing patch has been removed and greater than 10% crown cover remains. These areas remain in the woody extent.

The three classes are all reported as clearing activity but distinguished in ongoing SLATS annual reports of clearing areas. The full list of attribution pixel codes used during manual mapping and attribution of vegetation clearing is detailed in Table 5 together with how each informs both reporting of clearing and woody extent updating.

Table 5: Codes used in mapping and attribution of vegetation clearing, including landcover replacement class, partial, full attribution, and how each code informs reporting and updating of the woody extent.

	Code	Description	Examples	Reported as Clearing	Remain as Woody Vegetation
Human-induced clearing activity	40	Possible clearing	For review	No	Yes
	41	Full clearing to <i>pasture</i> : where vegetation has been converted from woody (i.e. >10% crown cover) to non-woody (i.e. <10% crown cover)	All clearing to <i>pasture</i> , internal property roads and tracks, fence lines, fire breaks. Single house pads in rural and semi-rural areas, golf courses	Yes	No
	71	Partial clearing major to <i>pasture</i> : where >=50% area of vegetation removed within patch and >10% crown cover remains	Clearing to <i>pasture</i> and still woody. Poisoning where dead trees left standing.	Yes	Yes

	73	Partial clearing major to <i>forestry</i> : where $\geq 50\%$ area of vegetation removed within patch and $>10\%$ crown cover remains	Clearing to <i>forestry</i> and still woody. <i>Forestry</i> includes timber harvesting on state- or privately-owned lands, where it can be verified.	Yes	Yes
	75	Partial clearing minor: where $<50\%$ area of vegetation removed and $>10\%$ crown cover remains	Disturbance including removal of understorey.	Yes	Yes
	42	Full clearing to <i>crop</i>	Within or adjoining existing cropping area where paddock has been ploughed for planting orchard/crops. Clearing of woody orchards (including grape vines) for horticultural purposes.	Yes	No
	46	Full clearing to <i>settlement</i>	Clearing for housing estates, shops, hospitals etc.	Yes	No
	48	Full clearing to <i>mine</i>	Mining activities including expansion of open cut mining, CSG pads, coal exploration areas and large quarries	Yes	No
	51	Full clearing to <i>infrastructure</i>	Dedicated roads, gas pipelines, rail & easements, airport runways and gravel pits beside roads. Dams and aquaculture ponds. Permanent linear features cleared between CSG pads.	Yes	No
	53	Full clearing to <i>forestry</i>	Clearing within State forests or exotic and native plantations.	Yes	No
	58	<i>Adjustment</i> : adjustment to the woody extent due to missed clearing	Correction for missed clearing in previous monitoring period.	No [†]	No
	59	<i>Adjustment</i> : adjustment to woody extent due to previous incorrect mapping	Correction to remove vegetation due to incorrect mapping.	No [†]	No
	80	<i>Intermediate code</i> : designates strip clearing. These are re-evaluated during post-processing and assigned either 41, 71 or 75). Refer to Section 3.3.5.4 for details.	Strip clearing and fodder harvesting but only in the Mulga lands.	Yes	Depends on final attribution.
Non-human-induced change	12	Fire	Active or recent fire scars	No	Yes
	16	Drying	Vegetation drier than in prev. year	No	Yes
	52	Flood damage	Change due to flooding event	No	**Yes
	54	Landslip damage	Change due to a landslip	No	**No
	56	Cyclone damage	Change due to cyclone	No	**Yes
	57	Natural tree death	Tree dieback where it can be identified and verified	No	**Yes

**Evaluated and attributed on case-by-case basis

[†]Reported as adjustment to woody extent, but not included in clearing areas.

3.3.5.3 Manual mapping methods

The manual mapping stage is critically important to production of the final high-quality woody vegetation clearing maps. The need for extensive manual editing is required because naturally occurring events can affect vegetation

in ways that appear similar to woody vegetation clearing in the imagery and clearing index. For example, damage by storms, fire and drought can all cause a reduction in canopy health or cover that can influence the woody vegetation clearing index in a similar way to a clearing event.

The manual mapping stage is conducted in two passes: initial editing of the automatically generated clearing classifications, followed by extensive review and quality assessment. For each mapping pass, automated processes (python scripts) recall the required image inputs to the analyst's local working directory. For a given Sentinel-2 tile this includes: the selected Sentinel-2 start and end date imagery, the automatically generated change classification and clearing index, as well as historical clearing data and any ancillary imagery used to aid interpretation. Editing/quality review is undertaken directly on the change classification raster. The woody vegetation clearing index (Section 3.3.4) provides a good starting point for the change classification, helping to focus the manual mapping effort on those areas where clearing is most likely to have occurred.

After each mapping pass, the edited change classification is versioned and archived and each editing stage is automatically logged in a spatial database table together with operator name and time stamps. All procedures are documented on the RSS Wiki.

All manual editing is supported by a set of documented mapping guidelines, which guides scientists in determining the full extent of the clearing footprint. This is a 'living' document, updated as procedures are refined, or decisions clarified. The manual editing work is done using *ERDAS IMAGINE* software which includes a suite of tools for viewing and enhancing imagery. The software allows editors to recode pixels in the change classification, enabling direct recoding of the change areas using the codes given in Table 5.

The first pass involves a scientist reviewing the change classification layer and associated Sentinel-2 imagery and identifying and recoding pixels in accordance with the codes in Table 5 and the Sentinel-2 mapping guidelines to map the full extent of the clearing event (i.e. the clearing footprint). Scientists draw upon their own expert landscape knowledge and that of others in the SLATS team to aid decision-making. Sometimes the pre-defined probability thresholds on the clearing index do not enable the clearing footprint to be fully mapped. In this case, pixels mapped as 'no change' are recoded and included in the clearing event. Some additional research was undertaken to try and further optimise these thresholds but was unable to yield any marked improvement. It should be noted that during the first pass, the scientists make their assessment guided primarily by the Sentinel-2 imagery with minimal reference to ancillary data to maintain consistency in scale and interpretation.

The second pass of the manual editing is undertaken by a senior DES remote sensing scientist to provide an independent check. It effectively replicates the first pass with some areas recoded, or areas which have been recoded in the first pass, highlighted for further review and discussion, where required. The mapping team regularly discuss issues, areas of interest and mapping specifications to ensure consistency and continuous improvement. During this second pass, the senior scientist may make judicious use of other high-resolution imagery sources to confirm difficult to interpret areas such as occurs where the vegetation is sparse, or the clearing activity is difficult to distinguish. The scientists make the required updates, and a script is run to save the final change raster to the HPC image archive.

This two-pass process ensures a high level of accuracy and consistency in the final map. Note that some western tiles, where there is historically very little or no clearing activity, are only subject to the first pass. Any areas in these regions that are uncertain during this single pass are discussed with senior scientists to maintain consistency.

3.3.5.4 Post-processing strip clearing mapping for the Mulga Lands bioregion (code 80)

Strip clearing and other fodder harvesting methods can result in complex clearing patterns which are difficult to map. This type of clearing activity is particularly prevalent in the Mulga Lands bioregion where fodder harvesting practices are common and some provisions in the VMA allow for this type of clearing to ensure stock can be fed during drought. Clearing patterns in this bioregion can include obvious strips which may exceed the SLATS specifications for linear features (i.e. 20m width) or complex criss-crossing patterns or narrow strips which are below the specifications for linear features. The SLATS change algorithm, while accurate at detecting strip clearing events, does not always provide enough detail to accurately delineate these various types of strip clearing without significant additional manual updating. Therefore, to ensure objectivity and repeatability, as well as accurate delineation of strip clearing in the Mulga Lands bioregion, a *post hoc* mapping process was developed.

During the first pass SLATS clearing mapping, strip clearing events that are not well-defined in the clearing index, or are difficult to efficiently map to SLATS specifications, are mapped by delineating the outer boundary of the clearing event and attributing the area as code 80 (an intermediate code). A *post hoc* process then applies an image segmentation model, trained by accurate mapping of woody vegetation in the local region, to provide an improved delineation of the cleared strips. This is then manually reviewed and refined as required to provide an accurate delineation of the clearing. The clearing is then attributed accordingly as either full or partial (major)

clearing and coded accordingly (i.e. codes 41 and 71 respectively; Table 5).

3.3.5.5 Post-processing the clearing mapping

What is effectively a third mapping stage is undertaken by the senior DES remote sensing scientist which moderates the clearing mapping across tiles. This involves some targeted, and some random, selection of tiles and moderating the clearing for accuracy and consistency. Changes and updates are made based on expert opinion, as required.

The clearing mapping for the individual Sentinel-2 tiles are then mosaiced into a single state-wide raster. An additional manual refinement exercise is undertaken to edge match the mapping in overlapping regions between adjacent images to ensure a seamless data set for the state.

The raster mosaic is polygonised using an ArcPy script.

Prior to integration into the woody extent master database, the clearing polygons are intersected with a forestry vector data set to ensure that clearing events that fall within regions where there are known forestry plantations are correctly attributed as *Clearing to forestry*. The forestry vector is derived from the union of *Agricultural land audit - current forestry plantations – Queensland* (Department of Agriculture and Fisheries) and *Queensland Digital Cadastral Database* (Department of Resources), where base tenure is *state forest*, *forest reserve* or *timber reserve*. These data sets are downloaded from the Queensland Government Spatial Catalogue for the date closest to the start of the given monitoring period.

The clearing polygons are then integrated into the woody extent geodatabase following the procedures of Section 4.1.1.2, as part of the annual woody extent updating.

3.3.6 Data products

The final SLATS state-wide clearing data set is stored in raster form, for use in the reporting framework, and vector form, for publication.

3.4 Woody vegetation age since last disturbance estimation

3.4.1 Overview

The initial aim was to estimate the age of woody vegetation in the 2018 baseline year, to then enable tracking of vegetation age over time in successive annual updates to the woody extent (refer Section 4.1.1) to provide environmental and contextual data that makes the woody extent, clearing and regrowth mapping more useful for a range of policy and land management purposes.

This research has resulted in a new approach to modelling land cover over long periods using a sampling framework that requires minimal temporal coverage. The method combines Landsat time series, the high-resolution woody extent, and a sequential classifier to estimate the likelihood of woody cover for each year over the length of the Landsat time series (30+ years).

Due to the limits of the time series-based age estimation, woody vegetation age is defined as: the estimated time since the woody vegetation was last significantly disturbed, or where possible to detect, the time since it started to grow, or regrow following a significant disturbance. Disturbance may be due to a clearing event or other disturbance such as fire, flood, drought-related tree death etc.

For a given Landsat pixel, the sequential classifier generates a 30+ year time series of woody probabilities. Heuristic rules are then used to detect patterns in the woody probability sequence that might characterise a typical persistent woody or regrowth response curve and track that behaviour over time. The estimated age from the probability sequence is combined with the SLATS Landsat clearing record to provide a preliminary estimation of age since last disturbance for all woody pixels recorded in the 2018 baseline woody extent and to track the age in subsequent woody extent updates.

3.4.2 Data

A thirty-three-year time series (1988 to 2021) of seasonal Landsat fractional cover components at 30m spatial resolution (Scarth *et al.*, 2008) was used to generate the revised 2018, 2019 and 2020 age since disturbance rasters. The fractional cover data consist of bare, green, and non-green sub-pixel components. Seasonal data were generated by calculating the medoid (Flood, 2013) for every three-month period (i.e. calendar season). Thus, there were 15 fractional cover components for every 12-month period (covering five seasons, Spring to Spring inclusive).

The 2019 woody extent data set was used as sample strata to generate woody/non-woody training sequences for a

Conditional Random Field classifier (described below in Section 3.4.3.1). The 2019 woody extent data were resampled from 10m to 30m spatial resolution to align spatially with the Landsat time series.

The historical SLATS Landsat (1988-2018) and Sentinel-2 (2018-2020) clearing data sets were additional inputs used to constrain the age since disturbance estimates for 2018, 2019 and 2020 woody extents.

3.4.3 Methods

3.4.3.1 Conditional Random Fields

Conditional Random Fields (CRFs) are a discriminative supervised classifier designed for sequential data (Lafferty *et al.*, 2001). CRFs are commonly used in the field of Natural Language Processing (NLP), which uses machine learning to understand sequential context in human languages. In the NLP field, a sequence might be represented by a sentence, and each word would consist of a set of predictive features. The goal of CRFs is to estimate the likelihood of transitioning between states (i.e. from one word to another). In this work, we used CRFs in a similar manner but applied to remotely sensed data. Therefore, our sequence for an individual pixel was a time series of seasonal fractional cover estimates. The sequence states were 12-month periods and the predictive features for each state consisted of fractional cover components over each 12-month timeframe.

3.4.3.2 Sample framework using temporal augmentation

CRFs are a supervised classifier and, therefore, require example class labels to optimise the state transitions. Collecting repeat training data for these types of temporal sequences over large areas is challenging. Therefore, a sampling method was developed that uses available land cover classifications over any number of time periods to generate sequences of training labels. This method is referred to as 'temporal augmentation'.

With this approach, the 2019 woody extent was used to generate random stratified samples over space, resulting in thousands of samples that intersected the manually scrutinised woody extent data set. In practice, further stratified samples with additional land cover edge stratum (e.g. woody cover 90 m from a non-woody pixel) were included to ensure that mixed pixels were represented in the sample pool. This random sample pool provided a starting point for woody/non-woody training examples. Next, training sequences (augmentation) from this static pool were generated.

The Landsat time series used for prediction were sequences of fractional cover over 33 years, but the training sample pool only included labels from the 2019 woody extent, which coincided with a single year of seasonal medoid fractional cover data (approximated as Spring 2018 - Spring 2019 incl.). Therefore, the samples were augmented to match the 33-year Landsat sequence length to train a CRF classifier. To do this, a subset of samples was selected within a mapping zone. A mapping zone could theoretically consist of any region, such as an ecological biome or administrative boundary. In this work, a regular grid of 150km x 150km across the state was used, with each grid cell representing an independent mapping zone. After woody/non-woody samples were collected for a particular mapping zone, thousands of 33-year pseudo-sequences were generated, where each pseudo-sequence consisted of samples spatially distributed across the mapping zone. It is important to note that each sample's class label in each sequence was always *woody* or *not woody* in the 2019 woody extent, and the associated Landsat data was always the set of seasonal cover fractions from the single year approximated by Spring 2018 - Spring 2019 (incl.). Therefore, these sequences were pseudo sequences because they did not represent real time series. By generating thousands of random permutations, though, this temporal augmentation approach provided a sequential data set for a CRF model to "learn" transitions between different land cover states (e.g., woody to woody, woody to non-woody, and non-woody to woody).

3.4.3.3 Woody probability estimates

The CRF classifier optimises the conditional distribution $p(y|X)$, which is the probability of y (e.g., a sequence of class labels) given a set of predictive features X (e.g., a sequence of predictive features). In this work, X and y were Landsat fractional cover time series and woody/non-woody label sequences, respectively, both taken from the temporal augmentation sample pool described in Section 3.4.3.2. More specifically, the predictive features consisted of 33 years of fractional cover sequences $[(bare, green, non-green)_{1988}, \dots, (bare, green, non-green)_{2021}]$, with corresponding woody/non-woody class label pseudo-sequences (e.g., $[woody_{1988}, \dots, non-woody_{2021}]$). These data were used to train CRF models for each 150km X 150km mapping zone. As described above, a CRF estimates the transition likelihood between land cover states. Thus, for an individual pixel the transition likelihoods between sequence states are estimated with the assumption that there is co-dependency between labels. For example, a CRF trained on the woody/non-woody sequential data estimates likelihoods as $[(p(w), p(nw))_{1988}, \dots, (p(w), p(nw))_{2021}]$, where the 1988 estimates might consist of a 0.7 and 0.3 likelihood of the pixel being predominantly woody and non-woody, respectively i.e. $(p(w)=0.7, p(nw)=0.3)_{1988}$.

3.4.3.4 Woody age since disturbance estimates

The start and end years of the *most recent period of regrowth or persistent woody* were estimated from the 1988-2021 time series of woody probabilities using a heuristic approach. Four thresholds guided this work: *minimum baseline probability threshold*, *minimum woody probability*, *minimum proportional change*, and *minimum number of consecutive years*. For production of the 2018, 2019 and 2020 age estimates, these thresholds were set to: 0.05, 0.5, 0.01 and 3, respectively.

For each pixel, the woody probabilities were evaluated to determine if the sequence at each time step was:

- a) *increasing*, characterised by a proportional change in probabilities exceeding the *minimum proportional change* and the probability exceeding the *baseline probability threshold*, or
- b) *woody*, characterised by the probability exceeding the *minimum woody probability threshold*. If the *increasing or woody* condition was met, the age was incremented.

If at a given time step, the incremental age exceeded the *minimum number of consecutive years as increasing or woody*, then the start and end years of that period were recorded. The start year recorded the time step at which the *increasing or woody* condition was first satisfied. The end year was incremented at each successive time step until either the end of the time series or until neither *woody* nor *increasing* conditions were satisfied, in which case, the incremental age was reset to zero, and the iteration restarted from that time step.

The SLATS historic Landsat (30m; 1988-2018) and Sentinel-2 (10m; 2018+) clearing data sets were used to constrain the modelled age estimates, so that the woody age since disturbance could not be greater than the time since last (full) clearing event recorded by SLATS:

- Where an historical SLATS clearing event was recorded for a given pixel, the age was estimated as the most recent of the last clearing event, and the estimated start of the last predicted regrowth/woody period.
- Where there were persistently high woody probabilities (exceeding *minimum woody probability*) over the entire sequence, an age equal to the length of the time series was assigned.
- Where no age could be estimated with any confidence due to an absence of clearing history and no detectable trends in the woody probability sequence, a label of “indeterminant” was assigned.

The Sentinel-2 partial clearing classes, and previous Landsat “Thinning” classes were excluded from use in age estimation.

Updates to the heuristic rules since the publication of *Statewide Landcover and Trees Study (SLATS) Sentinel-2 – 2018* have resulted in improved tracking of persistent woody vegetation over time, and increased ability to detect young regrowth. This in turn has resulted in fewer woody pixels with an ‘indeterminant’ age.

3.4.4 Limitations

There are some limitations to the woody vegetation age since disturbance estimation:

- The 2018 baseline woody extent contains sparse and/or young woody vegetation that may not have been detectable at the Landsat scale – either in the CRF modelling approach or by SLATS in previous monitoring periods and was therefore unable to be monitored at that scale.
- Missing data in the CRF fractional cover model inputs due to persistent cloud or topographic shadow as well as real but transient disturbance events such as fire or flood may cause woody probabilities to fall to 0, confounding age estimates.
- There may be clearing omissions from the SLATS record due to relative scale effects or mapping errors that impact the ability to estimate age for a given pixel.
- There may also be clearing events in the Landsat record that did not involve a full transition from woody to non-woody (according to 2018 woody baseline mapping specifications).
- Mismatches in scale and mapping specifications between the historic Landsat clearing data sets and the higher resolution baseline woody extent can result in artefacts in the age data set e.g. edge effects, small non-woody features such as roads and buildings which were the result of a past mapped clearing event but are currently included in the woody extent due to minimum mapping unit specifications.

3.4.5 Data products and outputs

Thirty-three years (1988-2021) of woody probabilities across the state at Landsat 30m spatial resolution have been produced. The data values indicate the likelihood (range of 0 to 1) that a pixel is predominantly woody.

The Queensland mosaic of estimated regrowth periods (1988-2021) captures the start and end year of the most recent detected regrowth or persistent woody period for each pixel over that period at Landsat 30m spatial resolution.

The woody age since disturbance rasters record the estimated age since disturbance (years) of woody pixels for the given year of woody extent at Sentinel-2 10m spatial resolution. Currently this includes 2018, 2019 and 2020 mosaics over Queensland with the intention to produce updates for each annual monitoring period.

3.5 Woody vegetation regrowth mapping

3.5.1 Overview

The aim is to monitor regrowth annually using Sentinel-2 imagery as part of the SLATS annual analysis of woody vegetation change, commencing with the 2019-20 monitoring period. In the first Sentinel-2-based monitoring period, 2018-19, it was assumed that existing regrowth has already been mapped and characterised in the 2018 baseline woody vegetation extent. As regrowth generally occurs on time scales greater than one year, mapping and monitoring of new regrowth is included in the 2019-20 monitoring period and will be included in annual monitoring thereafter.

Regrowth is only considered in the non-woody portion of the Queensland landscape as defined by the woody extent data set at the start of the given monitoring period. As such, densification of woody landscapes due to regrowth in already woody areas will not be considered at this stage.

The woody vegetation regrowth mapping approach is similar to that used for the clearing mapping in that a two-stage approach is used. Firstly, the automatic detection of candidate regrowth areas for consideration is undertaken using a similar CRF modelling approach as used for Landsat-based age estimation (Section 3.4) but applied to a shorter, rolling Sentinel-2 time series. This is then followed by a manual assessment and mapping process to determine if a candidate regrowth area is sufficiently woody to be counted as regrowth and added back to the woody account.

Where it can be mapped using the methods described below, regrowth will be delineated and added into the woody extent, effectively a 'gain' in the context of the accounting framework.

3.5.2 Methods

Following the work described in Section 3.4, time sequences of woody probabilities are used to predict candidate new regrowth areas. However, the regrowth monitoring framework differs from the age attribution in two key ways:

- i. a shorter, 10-year time series of Sentinel-2 imagery supplemented with up-sampled Landsat imagery is used instead of 30+ year Landsat sequence; and
- ii. regrowth is estimated with a modified set of heuristic rules adapted for the shorter time series and with a focus on young regrowth that is not currently in the woody extent.

3.5.2.1 Short-term woody probabilities

Sequential-based CRF woody probability estimates over ten years are utilised for this modelling approach. The timeframe chosen was designed to shorten the prediction sequence compared to the 33-year age estimates while maintaining sufficient temporal resolution to model temporal co-dependencies between woody/non-woody transitions. Any 10-year sequence ending around 2020 precedes the first Sentinel-2 launch date of 23 June 2015. Therefore, the Sentinel-2 time series is supplemented with Landsat data prior to the beginning of the Sentinel-2 era. Alternatively, the monitoring timeframe could be shortened to less than 10 years to use Sentinel-2 only. However, tests showed that sequences of less than five years were not sufficiently long enough for a CRF model to optimise land cover transitions. The Landsat is resampled from 30m to 10m spatial resolution to align with the Sentinel-2 grids. The practice of up-sampling coarser resolution imagery without ancillary information (e.g., fusion) is typically discouraged. However, in this case the Landsat data helped serve as a run-up period for the CRF model. That is, in practice, we are only interested in the final year of the 10-year period (refer Section 3.5.2.2) and only using the 10-year Landsat/Sentinel-2 sequence to help stabilise woody/non-woody transition estimates. The same 'temporal augmentation' approach is used to sample and fit a CRF model, using 10-year sequences instead of 33-year sequences.

As an example of how this works in the SLATS monitoring framework, for the monitoring period 2019-20, the 10-year time series consisted of resampled Landsat data from 2010 to June 2015. The remaining five years of the time series (i.e. 2015-2020) consisted of Sentinel-2 imagery only. For a given pixel, the sequence of woody likelihood from 2010-2020 (i.e. 10 annual probabilities) was estimated. These woody/non-woody probability sequences were then used in the heuristic model (refer Section 3.5.2.2) to estimate the presence (or not) of potential regrowth in 2020. This process was repeated for the following monitoring year (2021) by shifting the input time series to 2011-2021 and applying the same CRF model. The time series would ultimately consist of Sentinel-2-only during the 2025 monitoring year with a 2015-2025 time series. This approach could also be applied to timesteps of finer temporal granularity (e.g. monthly instead of seasonal input data), which would allow for a sub-annual monitoring

framework.

3.5.2.2 Regrowth candidate identification

Candidates for regrowth areas are identified using the rolling ten-year time series of woody probabilities (Section 3.5.2.1), and a heuristic classification similar to that used for woody age since disturbance estimation (Section 3.4) and with a similar but reduced set of thresholds: *minimum woody probability*, *minimum proportional change*, and *nconsec years* (the minimum consecutive number of years of regrowing or woody period). For the 2019-20 monitoring period these thresholds were: 0.5, 0.05, and 3, respectively.

As an initial condition, only pixels which are not already woody at the start of the given monitoring period are considered as potential candidates. For these non-woody pixels, the sequential woody probabilities are assessed to determine if a regrowth response is evident. Unlike in the age estimation, it is only the last period of *nconsec years* up to and including the end year of the monitoring period which is of interest. If the change in probabilities at each time step exceeds the *minimum proportional change* over the last *nconsec years* and the woody probability at the end year exceeds the *minimum woody probability threshold*, then the pixel is identified as an initial candidate.

As an example, for the monitoring period 2019–20, pixels which were non-woody in the 2019 woody extent data set were assessed to determine if they displayed a proportional increase in probabilities greater than 0.05 relative to the previous year, over all three years, 2018 – 2020 (inclusive). Those which met that condition and also had a woody probability exceeding 0.5 at 2020 were potential candidates.

Filtering was applied to the set of candidate regrowth pixels to remove clumps of candidate pixels smaller than 0.5ha.

3.5.2.3 Regrowth mapping

The mapping approach uses the automated regrowth predictions described in Section 3.5.2.2 above as candidate areas for regrowth which are each then manually assessed.

The manual process follows a similar structure to the clearing mapping described in section 3.3.5.3. Automated processes (Python scripts) are applied to a given Sentinel-2 tile to recall the Sentinel-2 imagery, the candidate regrowth estimates, and additional ancillary data for that tile into a local GIS (ArcGIS Pro). The ancillary data is used to aid in the decision-making process and includes historical clearing data, land use information, and additional high-resolution imagery when available. An initial editing stage using the CRF predictions is followed by quality assessment. Due to the often subtle and gradual increase in identifiable regrowth vegetation from satellite imagery, a time series of Sentinel images from 2016 through to 2020 was used by analysts to assess each candidate. The non-woody areas of each scene were scanned to ensure obvious regrowth was not missed by the CRF predictions.

Ideally regrowth needs to be observable in at least two consecutive years with obvious trees crowns identifiable in the imagery to be considered and mapped as regrowth. This ensures that non-woody ground cover is not incorrectly included in the mapping product. Identified regrowth is classified using the attributes listed in Table 6. As with the clearing mapping, these attributes are intended to provide an indication of the purpose for which the land is used at the location where the regrowth is mapped. Where potential regrowth is identified but has not developed to a suitable stage, or uncertainty exists, the area will be flagged for potential regrowth to be assessed again in the following mapping period. Quality assessment is conducted by a Senior Remote Sensing Scientist to check and validate mapped regrowth.

Field work has been undertaken to validate that mapped regrowth in the given monitoring period is woody ($\geq 10\%$ crown cover), and to confirm that the correct code had been applied. Where possible, features mapped as “Possible regrowth” are prioritised for field observations to improve certainty in the mapping, calibrate mapping staff in the image interpretation of woody vegetation and improve landscape knowledge. Field observations of young woody regrowth which is not yet observable in Sentinel-2 imagery are captured opportunistically. These areas are then monitored for future regrowth mapping.

Table 6: Codes used for the SLATS regrowth mapping attribution.

	Code	Description	Examples
Regrowth Activity	3	Possible regrowth	Vegetation that has been identified as possible regrowth, but there is insufficient evidence to include it in the reporting. Flagged for future monitoring.
	21	Regrowth in <i>Pasture</i>	All native and non-native woody vegetation regrowth in <i>pasture lands</i> .
	22	Regrowth in <i>Cropping</i>	New and regrowing woody crops and horticulture, including fodder crops and tree fruits and nuts.
	23	Regrowth in <i>Forestry</i>	Regrowth in timber plantations and state forests. <i>Forestry</i> includes timber on state- or privately-owned lands, where it can be verified.
	24	Regrowth in <i>Mine</i>	Regrowth in mining areas. i.e., rehabilitation.
	26	Regrowth in <i>Settlement</i>	Regrowth in urban residential areas such as gardens and parklands.
	27	Regrowth in <i>Infrastructure</i>	Regrowth along rail lines, airports, industrial complexes.
	28	<i>Adjustment: incorrect clearing</i>	Incorrect full clearing in the previous mapping period added back to the woody extent. Reported as an adjustment.
	29	<i>Adjustment: incorrect non-woody vegetation</i>	Correction to add vegetation due to incorrect mapping in the woody extent. Reported as an adjustment.

3.5.3 Data products and outputs

After finalising, the regrowth together with the clearing data sets are integrated with the woody extent data set to provide an updated woody extent and change attributes (Section 4.1.1).

The final SLATS state-wide regrowth data set is stored in raster form, for use in the reporting framework, and vector form, for publication.

3.6 Woody vegetation density estimation using foliage projective cover

3.6.1 Overview

Foliage projective cover (FPC) is a metric of vegetation cover used in many Australian vegetation classification frameworks. In the current program, SLATS uses the FPC metric derived from Sentinel-2 imagery to provide broad estimates about the range of tree and shrub densities represented in woody vegetation across Queensland, and in the context of the monitoring framework, the vegetation clearing and regrowth mapping.

For some time, RSS has produced FPC using a model applied to Landsat, calibrated by field observations (Armston, 2009, Kitchen, 2010). An updated model was developed which relates field measurements of FPC to 2-year time series of NDVI computed from Landsat seasonal surface reflectance composites. As with the woody vegetation clearing index (Section 3.3.3.2), there is insufficient field data that coincides with the Sentinel-2 satellite program. The model was therefore fitted on Landsat imagery using a significantly expanded set of field data than was used for the previous Landsat FPC model fitting. The model is then applied to analogous Sentinel-2 seasonal surface reflectance composites to produce an FPC image, using the radiometric relationships established between Sentinel-2 and Landsat in Flood (2017). This is intended to represent the FPC for that 2-year period rather than any single date.

The data set is generally expected to provide a reasonable estimate of the range of FPC values for any given stand of woody vegetation, but it is expected there will be over- and under-estimation of absolute FPC values for any specific location (i.e. pixel).

3.6.2 Data

The updated FPC model was developed based on:

- A two-year (eight- season) time series of Landsat seasonal surface reflectance composites (using medoid method of Flood, 2013b). Use of the seasonal composites aims to reduce noise and cloud contamination in

surface reflectance while still capturing much of the seasonal variation.

- The national data set of historic star-transect field data (Muir *et al.*, 2011) from which measures of FPC can be derived. At the time of the model development, this was in excess of 4000 individual sites.

In the model, FPC is the combined green fraction of foliage from over- and mid-storey foliage from woody plants (trees and shrubs). While it would be desirable to have a model fitted to Sentinel-2 imagery, little of the available field data has been collected since the launch of Sentinel-2.

To produce a Sentinel-2 FPC product that represents a given period, the model is applied to two-year time series of Sentinel-2 seasonal surface reflectance composites, radiometrically adjusted to match Landsat using relationships described in Flood (2017).

For SLATS monitoring and reporting, the two-year time series for FPC production is selected to best estimate FPC of woody pixels for a given woody extent data set. The start and end seasons are chosen to best approximate the SLATS annual monitoring period (nominally August through to August) and so that *all* seasonal input data precede the nominal date of woody extent (August of given year). This ensures that the maximum NDVI pixel used to estimate FPC for the given woody extent is not influenced by woody vegetation change that occurs in the successive monitoring period. As an example, the published *Statewide Landcover And Trees Study (SLATS) Sentinel-2 – 2018 Foliage Projective Cover (FPC) – Queensland*, estimates FPC for the 2018 woody extent, using season reflectance composites from the two years preceding: Winter 2016 – Autumn 2018 (inclusive).

3.6.3 Methods

3.6.3.1 Model fitting

The FPC model relates the field measured green fraction of mid- and over-storey foliage cover to the minimum value of NDVI calculated from 2-years of Landsat seasonal surface reflectance composites. NDVI is a standard vegetation index used in remote sensing which is highly correlated with vegetation photosynthesis (Rouse, 1974). Other indices and metrics were tested but yielded equivalent or worse results therefore NDVI was chosen for simplicity and as it is a widely known index of vegetation cover.

The FPC model is sensitive to fluctuations in vegetation greenness, leading to anomalies such as high FPC on irrigated pastures or locations with very green herbaceous or grass understoreys. A given pixel in the FPC image, represents the predicted FPC in the season with the least green/driest vegetation cover over the 2-year period assumed to be that with the least influence of seasonally variable herbaceous vegetation and grasses on the more seasonally stable woody FPC estimates. The two-year period was used partly because it represents a period relative to tree growth but was also constrained due to the limited availability of imagery in the early Sentinel-2 time series. Other time periods were tested but did not improve estimates.

The fitted model is a simple quadratic model with three coefficients.

$$FPC = C_0 + C_1NDVI_{\min} + C_2NDVI_{\min}^2$$

where $NDVI_{\min}$ is the second lowest value of NDVI for the 2-year period of seasonal surface reflectance and C_0 , C_1 , C_2 are the model coefficients. The second lowest value was chosen as cloud contamination due to failure of cloud masking in the seasonal surface reflectance composites, may also result in very low values of NDVI.

Model performance was assessed with Monte Carlo Cross Validation (MCCV).

3.6.3.2 FPC prediction

The FPC model is applied to the two-year time series of Sentinel-2 seasonal surface reflectance composites to produce an FPC image. Finally, the woody extent data set (Section 3.2.6) is used to reset FPC values in non-woody regions to zero to eliminate over-estimation of FPC in green pastures, cropping regions and other non-woody landscapes.

3.6.4 Data products and outputs

The final product is a state-wide Sentinel-2 Foliage Projective Cover estimated for a given woody extent and used in reporting of woody vegetation density. It is also released as a stand-alone data product as it is in high demand for a range of applications, particularly related to carbon, fire and biodiversity programs, and grazing (i.e. pasture biomass) modelling and prediction.

4 Data integration, reporting and accounting

4.1 Data integration

4.1.1 Annual updating of the woody extent

The woody extent (Section 3.2) forms the foundation of the SLATS woody vegetation (annual) change accounting framework (Section 2.2). In each annual monitoring period, once change mapping has been finalised, the woody extent is updated by integrating the SLATS clearing (from 2018-19 onward), and regrowth mapping (from 2019-20 onward). For a given monitoring period, both the woody extent and change are used in reporting. The updated woody extent is then used to initialise change mapping for the next period of mapping and reporting. As an example, in the 2019-20 monitoring period, the 2019 woody extent is updated using the 2019-20 clearing and regrowth to create a 2020 woody extent, which is then used as the basis for 2020-21 change mapping.

During the initial transition to the Sentinel-2-based monitoring (over 2018-19), while the woody extent data set and enhanced monitoring framework were still under development there were several transitional steps required to integrate the change mapping and woody extent data sets (Sections 4.1.1.1 and 4.1.1.2). Following this transitional period, annual updating routinely involves adding areas of regrowth to, and taking away areas of clearing, from the woody extent (Section 4.1.1.1). Areas of partial clearing (minor and major partial clearing activity) remain in the woody extent but are attributed for reporting purposes and to inform future monitoring.

The maintenance of the woody extent is not as straightforward as subtracting the clearing and adding the regrowth each year. It is inevitable that there will be ongoing refinements due to misclassification and error. Annual updating and maintenance of the woody extent allows for corrections due to errors in the woody extent, regrowth or clearing mapping in previous eras. Updates due to corrections are integrated similarly to clearing and regrowth but are coded and reported separately to distinguish additions and subtractions due to error/mapping refinement.

4.1.1.1 2018 update

The initial woody extent map (Section 3.2) was derived from 2017 Earth-I imagery. As a 2018 woody extent baseline was required for the revised SLATS accounting framework, the first step was to update the 2017 map to 2018 and establish the baseline.

The previously released 2017-18 SLATS Landsat clearing data set was used to inform the update of the woody extent data set from 2017 to the 2018. Differences in scale, specifications, and definitions between the 2017 woody extent and the 2017-18 Landsat-based clearing data set presented some difficulties for seamless integration of the two data sets. The SLATS 2017-18 clearing data set was produced using Landsat imagery, at a period prior to the incorporation of the woody extent data set into the SLATS clearing mapping process. As such, the resolution is coarser than the woody extent product. Further, the pixel-based Landsat clearing mapping approach differs from the current Sentinel-2 footprint mapping approach (refer to Section 3.3.5.1). The definition of “woody” also varied between the two data sets. The minimum woody density for inclusion in the woody extent is ~10% crown cover (~5% FPC) as mapped from Earth-i and Sentinel-2 imagery. The Landsat-based method used a more conservative woody threshold of ~20% crown cover (~10-11% FPC), or that which is detectable in Landsat imagery. Finally, the 2017-18 Landsat-based SLATS clearing data set did not include the partial clearing categories used in the current method (Section 3.3.5.2) which identify clearing activity where there has been significant modification, but enough woody material remains to be retained in the woody extent data set.

A comprehensive manual checking and editing approach was used to update the SLATS 2017-18 clearing data to align with the scale and definition of the 2017 woody extent to allow for easier integration. Scientists checked each clearing event greater than 2ha using Sentinel-2 imagery for the relevant monitoring end date. A Sentinel-2 based clearing index was also used to identify missed clearing not recorded in the 2017-18 Landsat clearing data set. Manual editing was applied to include missed clearing from the 2017-18 monitoring period, and to fully delineate the whole clearing footprint in line with the current approach for clearing mapping. Each event was also reviewed to determine if full or partial clearing had occurred as per the current approach. Full clearing areas were removed from the woody extent and partial clearing remained in the woody extent map. The updated woody extent was filtered to remove polygons <0.5ha, the minimum mapping unit used in baseline establishment. The result was a 2018 statewide baseline woody extent map.

4.1.1.2 2019 update

Integration of clearing mapping and woody extent for the first Sentinel-2-based 2018-19 monitoring period was more seamless, as the Sentinel-2 clearing and woody extent data sets are more integrated from the outset of change mapping, and more closely aligned in mapping scale and resolution. As the program was still refining

partial and full clearing mapping categories, there was some additional assessment and editing required to further refine these areas as part of the 2019 update. This was particularly the case in areas such as the Mulga Lands bioregion where strip clearing and other fodder harvesting management resulted in some complex clearing patterns and hence some difficulty determining full clearing (updated to non-woody in the woody extent) or partial clearing (remaining in the woody extent). Where cleared strips could be defined and met the woody extent criteria for linear features of 20m minimum width, they were manually delineated and coded as full or partial (major) clearing (codes 41 and 71 respectively, Table 5). Following post-processing attribution and filtering (described in Section 4.1.1.2) the result was a 2019 update to the statewide woody extent map.

4.1.1.1 Ongoing annual updates

From 2019-20 onwards, the SLATS methods are more fully established, and updates to the woody extent routinely incorporate annual gains due to regrowth as well as losses due to clearing. For example, the 2020 update involved subtracting the 2019-20 full clearing, adding the 2020 regrowth, attributing woody polygons impacted by partial clearing activity and integrating corrections for incorrect woody extent or previous clearing mapping. The development of an automated method for more refined mapping in complex woody vegetation clearing patterns associated with fodder harvesting and strip clearing specifically (Section 3.3.5.4) has also been finalised.

For the foreseeable future woody extent updates against the finalised change mapping will continue to be a post-hoc data integration exercise. The longer-term vision for SLATS is a fully integrated editing framework such that all editing is done directly on a master woody extent layer. Change attribution would be managed through the data set's attribute table and adding new polygons or splitting existing polygons.

4.1.1.2 Post-processing and attribution

A series of Arcpy batch scripts is used to process both the clearing (Section 3.3) and regrowth (Section 3.5) polygons, clip them into the woody extent data set, update the attribution, and finally to apply some cleaning and filtering to remove artefacts associated with the editing and updating procedures. Finally, the updated woody extent and change polygons are filtered to remove slivers and patches which are less than 0.25ha to ensure the ongoing monitoring specification of 0.25ha minimum mapping unit are maintained in the data set.

For a given monitoring period, a new attribute field is added to the master woody extent database to record the updated woody extent attribute for each change polygon. Another is added to record the clearing or regrowth attribution codes (Table 5 and Table 6 respectively) for that monitoring period. For example, upon completion of the 2019-20 change mapping, the clearing and regrowth polygons were incorporated into the master woody extent database by adding a *2020* field, where polygons were recorded as either woody or non-woody depending on the clearing or regrowth attribution codes and a *2019-20 Change* field was added to record the clearing and regrowth attribution codes. Full clearing and regrowth involve a conversion from woody to non-woody and non-woody to woody respectively. Partial clearing involves only an update in the Change field for the relevant period. All remaining unchanged woody extent polygons are replicated into the attribute field corresponding to the updated year. The result is a single data set containing multiple epochs with a separate field for each woody extent year and fields recording the landcover class attributed to the change for each period.

4.2 Reporting

4.2.1 The reporting package

The data products described in the sections above provide the basis for combining SLATS information with other data to monitor and account for woody vegetation extent and change in Queensland on an annual basis. Annual reports are released as web-based reports. The spatial data sets and annual reports are made available via the SLATS data products web page, with supporting data downloadable from the reporting pages.

This enables SLATS information and data to be accessible to a range of stakeholders and users in formats that service a range of requirements and analyses.

The SLATS reporting series associated with the current methodology commenced with 2018 baseline woody vegetation extent and 2018-19 clearing reports, and from 2019-20 onwards, annual reporting of clearing and regrowth.

4.2.1.1 The SLATS cross-tabulation and regional data summaries

The area of change in Queensland is summarised in cross-tabulation using several grouping layers including those listed in Section 4.2.1.3, but also including a wider range of regional and administrative boundaries:

- Local Government Areas
- Natural Resource Management regions
- Catchments
- Biogeographic subregions

This creates a large table, from which flexible summaries can be easily made using simple queries. For example, the area of woody vegetation for 2019 can be calculated by the sum of the area in 2018 less the area mapped as cleared. More complex breakdowns of clearing activity, for example, by Regulated Vegetation Management (RVM) Map category, catchment and clearing type can be summarised.

This change table forms the basis for the web-based reporting and the regional data summaries available as Open Data. It also provides a simple, pre-processed mechanism for arbitrary reporting on request.

The procedure for creating the table is captured in a dedicated git repository. The workflow makes use of the Snakemake package (Mölder et. Al. 2021) and container processing (Kurtzer, Sochat & Bauer 2017) to ensure repeatability and process governance.

4.2.1.2 The 2018 baseline woody vegetation extent report

The 2018 baseline woody vegetation extent report provides the initial baseline for ongoing SLATS reporting, which annually monitors and accounts for woody vegetation extent and change in Queensland. The revised SLATS methodology now monitors and report changes in woody vegetation extent against this baseline. The report also includes information relating to woody vegetation density and age since disturbance estimates.

4.2.1.3 The annual SLATS report

The annual SLATS report summarises, and reports changes due to gains (regrowth) and losses (clearing) in woody vegetation across Queensland for the nominal period from August to the following August. The first of these reports, for the 2018-19 monitoring period, only includes changes due to woody vegetation clearing activity—regrowth reporting is not included. It is assumed that existing regrowth in 2018-19 has been mapped and characterised in the 2018 woody vegetation extent baseline (Sections 3.5 and 4.2.1.2).

Annual SLATS reports provide the key findings and a range of graphical and tabulated woody vegetation change activity summarised by:

- regrowth and type of clearing activity: full, partial (major) and partial (minor) clearing
- landcover replacement class
- broad categories of age and density of woody vegetation
- vegetation management status including RVM Map category and vegetation management class
- bioregions, with other regional summaries available via open data e.g. Local Government Areas, catchments
- a transaction summary of woody vegetation change over the period.

4.2.1.4 Data products

Data products produced by the program are prepared and released as Open Data. This includes the clearing and regrowth data, the woody extent data, the age and density products, and any data associated with reporting and regional summaries.

4.2.1.5 A note on reporting areas vs rates

Historically, SLATS has reported annual rates of clearing mostly due to the difficulty of acquiring two cloud-free dates of Landsat imagery one year apart (Appendix A in Queensland Department of Environment and Science, 2018). For example, the start and end dates selected could be as little as 8 months, or as much as 15-16 months apart, sometimes for adjacent path/rows. Thus, clearing *areas* were adjusted (i.e. annualised) to account for this time lag and to enable comparative annual reporting. The increased temporal resolution of the Sentinel-2 satellites (5-day return interval vs Landsat's 16-day) means that opportunities for obtaining cloud-free imagery much closer to 12 months apart are significantly increased. For example, an analysis of the SLATS-selected cloud-free Sentinel-2 tiles for 2019 showed that if a nominal date of August 15 was chosen, all the Sentinel-2 tiles for the state were within one month of that date. The only tiles which were not within one month were some coastal tiles in the far North of Cape York which have very little landmass and are generally cloud-affected year-round. As such, the revised program will no longer report annualised rates, rather reporting the actual areal change figures. This generally makes reporting simpler and perhaps most importantly, simplifies the reporting and communication for a general audience.

5 References

- Ahmad, W., D.L.B. Jupp, and M. Nunez. 1989. "Use of Remotely Sensed Data for Land Cover Mapping in a Mountainous Region of North East Tasmania, Australia." In *Collected Conference Papers Describing Remote Sensing Application Projects Using MicroBRIAN Image Processing.*, 33–48.
- Armston, J., Denham, R., Scarth, P., Moffiet, T. 2009. Prediction and validation of foliage projective cover from Landsat-5 TM and Landsat-7 ETM+ imagery. *Journal of Applied Remote Sensing* 3 (1) 033540 033540 <https://doi.org/10.1117/1.3216031>
- European Space Agency, 2019, S2 MPI L1C Data Quality Report, Issue 46. 05/12/2019. Ref: S2-PDGS-MPC-DQR, ESA Copernicus. https://sentinel.esa.int/documents/247904/685211/Sentinel-2_L1C_Data_Quality_Report/
- Flood, N., Buck, R. and Teutsch, C. Unpublished. An Index of Woody Vegetation Clearing Using Sentinel-2 Imagery
- Flood, N. 2013a. An operational scheme for deriving standardised surface reflectance from Landsat TM/ETM+ and SPOT HRG imagery for Eastern Australia. *Remote Sensing*, 5 (10) pp 83-109
- Flood, N. 2013b. Seasonal composite Landsat TM/ETM+ images using the medoid (a multi-dimensional median). *Remote Sensing*, 5(12) pp6481-6500.
- Flood, N., Danaher, T. 2013c. A File Naming Convention for a Large Satellite Imagery Archive: QVF – a case study. Figshare. Journal contribution. <https://doi.org/10.6084/m9.figshare.875298.v1>
- Flood, N. 2017. Comparing Sentinel-2A and Landsat 7 and 8 using surface reflectance over Australia. *Remote Sensing*, 9 (7), 659.
- Flood, N., Watson, F. and Collett, L. 2019. Using a U-net convolutional neural network to map woody vegetation extent from high resolution satellite imagery across Queensland, Australia. *International Journal of Applied Earth Observation and Geoinformation*, Volume 82. <https://doi.org/10.1016/j.jag.2019.101897>.
- Flood, N., Danaher, T., Gill, T., Scarth, P. 2020. Forest Clearing with SPOT-5. figshare. Online resource. <https://doi.org/10.6084/m9.figshare.12980897.v1>
- Kitchen J, Armston J, Clark A, Danaher T, Scarth, P. 2010. Operational use of annual Landsat-5 TM and Landsat-7 ETM+ image time-series for mapping wooded extent and foliage projective cover in north-eastern Australia, in *Proceedings of the 15th Australasian Remote Sensing and Photogrammetry Conference*, Alice Springs, Australia, 13–17 September 2010.
- Kurtzer GM, Sochat V, Bauer MW. 2017. Singularity: Scientific containers for mobility of compute. *PLOS ONE* 12(5): e0177459. <https://doi.org/10.1371/journal.pone.0177459>
- Lafferty, J., McCallum, A., and Pereira, F CN. 2001. Conditional random fields: Probabilistic models for segmenting and labelling sequence data. University of Pennsylvania Engineering Departmental Papers.
- Mölder, F., Jablonski, K.P., Letcher, B., Hall, M.B., Tomkins-Tinch, C.H., Sochat, V., Forster, J., Lee, S., Twardziok, S.O., Kanitz, A., Wilm, A., Holtgrewe, M., Rahmann, S., Nahnsen, S., Köster, J. 2021. Sustainable data analysis with Snakemake. *F1000Res* 10, 33.
- Muir, J., Schmidt, M., Tindall, D., Trevithick, R., Scarth, P., & Stewart, J. 2011. Field measurement of fractional ground cover: a technical handbook supporting ground cover monitoring for Australia. Technical report, Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra, ACT.
- NSW Department of Planning, Industry and Environment. 2020. Summary woody vegetation change, Statewide Landcover and Trees Study (SLATS) for 2018. NSW Government, Sydney.
- Olofsson, P., Foody, G. M., Stehman, S. V., & Woodcock, C. E. 2013. Making better use of accuracy data in land change studies: Estimating accuracy and area and quantifying uncertainty using stratified estimation. *Remote Sensing of Environment*, 129, 122–131. <https://doi.org/10.1016/j.rse.2012.10.031>.
- Olofsson, P., Foody, G., Herold M., Stehman, S., Woodcock, C., Wulder, M. 2014. Good practices for estimating area and assessing accuracy of land change, *Remote Sensing of Environment*, 148, pp 42-57.
- Pickering, J., Stehman, S., Tyukavina, A., Potapov, P., Watt, P., Jantz, S., Bholanath, P., Hansen, M. &. 2019. Quantifying the trade-off between cost and precision in estimating area of forest loss and degradation using probability sampling in Guyana. *Remote Sensing of Environment*, 221 pp 122-135. <https://doi.org/10.1016/j.rse.2018.11.018>
- Queensland Herbarium. 2019. Remnant regional ecosystem vegetation in Queensland, Version 11.0, Department

of Environment and Science, Brisbane.

Queensland Land Use Mapping Program (QLUMP), Department of Environment and Science,
<https://www.qld.gov.au/environment/land/management/mapping/statewide-monitoring/qlump>

Queensland Department of Environment and Science. 2018. Statewide Landcover and Trees Study (SLATS): Overview of Methods. DES, Brisbane.

Ronneberger, O., Fischer, P., Brox, T. 2015. U-net: Convolutional networks for biomedical image segmentation. In: Navab, N., Hornegger, J., Wells, W.M., Frangi, A.F. (Eds.), *Medical Image Computing and Computer-Assisted Intervention – MICCAI 2015*. Springer International Publishing, Cham, pp. 234–241. https://doi.org/10.1007/978-3-319-24574-4_28.

Rouse, J.W., Haas, R.H., Scheel, J.A., and Deering, D.W. 1974. Monitoring Vegetation Systems in the Great Plains with ERTS. In *Proceedings, 3rd Earth Resource Technology Satellite (ERTS) Symposium*, vol. 1, p. 48-62. <https://ntrs.nasa.gov/citations/19740022614>

Scarth, P, S Gillingham, and J Muir. 2008. Assimilation of Spectral Information and Temporal History into a Statewide Woody Cover Change Classification. In *Proc of 14th Australasian Remote Sensing and Photogrammetry Conference*. Alice Springs, 2008

Scarth, P., Armston, J., Lucas, R., and Bunting, P. 2019. A Structural Classification of Australian Vegetation Using ICESat/GLAS, ALOS PALSAR, and Landsat Sensor Data. *Remote Sensing* 11, no. 2: 147. <https://doi.org/10.3390/rs11020147>

Specht, R. 1983. Foliage projective covers of overstorey and understorey strata of mature vegetation in Australia. *Austral Ecol.*, 8 (4), pp. 433-439. <https://doi.org/10.1111/j.1442-9993.1983.tb01340.x>.

Storey J, Choate M, and Lee K. 2014. Landsat 8 Operational Land Imager On-Orbit Geometric Calibration and Performance *Remote Sensing*, 6(11), pp. 11127–11152. <https://doi.org/10.3390/rs6111127>.

Strahler, A. H., Boschetti, L., Foody, G. M., Friedl, M. A., Hansen, M. C., Herold, M., *et al.* 2006. Global land cover validation: Recommendations for evaluation and accuracy assessment of global land cover maps. EUR 22156 EN – DG. Luxembourg: Office for Official Publications of the European Communities (48 pp.).

Zhang, W., M. Brandt, Q. Wang, A. V. Prishchepov, C.J. Tucker, Y. Li, H. Lyu, and R. Fensholt. 2019. From woody cover to woody canopies: How Sentinel-1 and Sentinel-2 data advance the mapping of woody plants in savannas. *Remote Sensing of Environment* 234:111465. <https://doi.org/10.1016/j.rse.2019.111465>.

Zhu, Z., Woodcock, C. 2012. Object-based cloud and cloud shadow detection in Landsat imagery. *Remote Sensing of Environment*, 118, pp. 83-94. <https://doi.org/10.1016/j.rse.2011.10.028>

Zhu, Z., Wang, S., Woodcock, C. 2015. Improvement and expansion of the Fmask algorithm: cloud, cloud shadow, and snow detection for Landsats 4–7, 8, and Sentinel 2 images. *Remote Sensing of Environment* (159) pp 269-277. <https://doi.org/10.1016/j.rse.2014.12.014>

6 Appendix

6.1 Landsat continuity: a sample-based estimation of woody clearing in 2018–19 using SLATS Landsat-based methodology

6.1.1 Introduction

SLATS has been detecting woody vegetation clearing based on Landsat for over 30 years. As detailed in the previous sections of this document, SLATS has undergone significant methodological changes starting from the 2018-19 monitoring period. In addition to increasing the spatial resolution from 30m to 10m by changing the satellite base imagery to Sentinel-2, the program has adopted a more comprehensive approach in which the clearing footprint is mapped as cleared instead of just single pixels of woody change. This treatment of clearing as a landscape phenomenon is supported by the incorporation of the woody vegetation extent product into the mapping process to identify which patches are woody. These methodological changes mean that clearing estimates obtained from this new approach are not comparable to previous clearing estimates based on Landsat.

The inability to compare the 2018-19 clearing results with previous monitoring periods is particularly problematic as it becomes difficult to assess the impact of legislative changes. A series of amendments to the VMA were passed in 2018 to reintroduce a number of the laws which were repealed by the previous government, and to repeal some laws which were contributing to increased clearing. The intention of the amendments was primarily to reduce clearing rates in regulated vegetation. However, the effect of the amendments would not be directly measurable because the estimates of clearing based on the revised SLATS methodology are not directly comparable with previously reported clearing rates. Furthermore, as the new methodology involves a change in the way that the clearing is being mapped, it is not possible to simply model the predicted Landsat-based area using a degradation model. Therefore, there is a need to generate estimates of clearing that are Landsat based and that follows the previous SLATS methodology.

The aim of this work is to present a sample-based estimate of the area cleared based on the SLATS Landsat-based methodology to provide an estimate of clearing in 2018-19 which can be compared with previously reported SLATS clearing rates. In other words, this work looked at what would have been mapped if SLATS had continued a similar mapping process based on Landsat.

6.1.2 Methods

Given time and resource limitations, it was not feasible to run a parallel Landsat/Sentinel-2 change mapping exercise. As an alternative, a sample-based approach in a subset of the state was developed. The established SLATS methodology for the mapping clearing was followed (Queensland Department of Environment and Science, 2018).

6.1.2.1 Study area

The study area was the Brigalow Belt Bioregion. This area was selected because it has historically been an area of relatively high clearing rates, has a range of land uses and ecosystems and includes over half of the Great Barrier Reef catchment area.

6.1.2.2 Satellite data

6.1.2.2.1 Landsat scenes

Landsat scenes from 2018 were compared to selected scenes from 2019. The 2018 scenes were those that had been previously used for the end date of the 2017-18 monitoring period. For 2019, 21 Landsat scenes that cover the Brigalow Belt Bioregion were selected using the following criteria:

- a) scene date should be as close as possible to the Sentinel 2 date for 2018-19;
- b) images should be from the dry season to maximize contrast, and
- c) scenes with the lowest cloud cover and similar green/dryness between Landsat and Sentinel-2 were preferred. No composites were used.

6.1.2.2.2 Clearing classification

The Landsat-based clearing index (Scarath *et al.*, 2008) was applied to the pairs of Landsat dates to produce a clearing probability layer which could then be edited by SLATS scientists.

6.1.2.2.3 Ancillary data

AIRBUS very high-resolution resolution imagery (<https://www.intelligence-airbusds.com/imagery/oneatlas/data/>)

and Sentinel 2 were used to guide decisions. Ancillary data is used to assist determinations for woody/non woody vegetation, and if clearing is identified, confirm that the changes are indeed human-induced clearing.

6.1.2.2.4 Sampling design

A grid that covers the Brigalow Belt Bioregion was produced using geoprocessing tools from ArcPro v2.4.2. The area was divided into 5,189 units (Figure 2). The size of each grid cell was defined based on the mapping scale used in the interpretation (1:37000). Each cell corresponds to 11.5km by 6.5km, which is the size of the window desktop viewer at the scale used by the editing scientist. The grid was overlaid with the 21 Landsat scene footprints, and information on the path/row was assigned to each cell along with a unique identifier. Each grid cell corresponds to one sampling unit.

A series of eligibility criteria were applied to the sampling units to filter those units that were considered impractical to include in the sample. To be included in the sample, units should be completely located within a scene and contain at least 70% of valid pixels (i.e. coastal units with only a small portion of land were removed). In addition, only units that contain at least one probability pixel derived from the clearing index classification were included. This criterion was based on analysis of previous monitoring periods which demonstrate that areas without an initial indication of potential clearing are very rarely identified as being cleared. A simple random sample of 20% of the eligible sampling units per scene was taken. To provide information on operator variability, a second sample of 20% was taken from 5 scenes, half of which had been evaluated in the first round.




Figure 2. Sample units for the study

6.1.2.3 Image interpretation process

Each unit in the sample was assessed using the same interpretation approach for SLATS as previous monitoring periods. To account for interpretation variability half of the sampling units in five of the scenes were interpreted twice by different scientists. The scientists undertaking the editing did not know if the unit that was being assessed

had been previously evaluated or not. As with standard SLATS methodology, each of the sampling units that was edited was checked by a senior scientist to maximize consistency and guarantee the quality of the product.

6.1.2.4 Sample-based estimation of the area cleared

The proportion of cleared pixels from all the valid pixels in each sampling unit was estimated. A cross tabulation between the edited raster and the most recent remnant vegetation map, for each sampled unit was done. Each unit was then summarised as the proportion of pixels marked as cleared relative to the total number of valid pixels (excluding cloud, no data, water etc pixels) in the unit.

The mean proportion of pixels that were cleared in remnant and non-remnant vegetation was estimated. The estimated area cleared was obtained as the product of the mean proportion of cleared pixels by the number of valid pixels within the Brigalow Belt Bioregion for 2017-18. Results were converted from pixels to ha by multiplying by 0.09 (each pixel 900m²).

6.1.3 Results

The estimated area of total clearing in the Brigalow Belt in 2018-19 was 234,810 hectares, an increase of 38,160 hectares compared to the 2017-18 period (Table 7). The 95% confidence interval suggests that this is a significant increase from the 2017-18 period

Table 7: Summary of estimated total clearing for the Brigalow Belt bioregion in the 2018-19 period

Veg Class	Area (1000's ha)			Change	
	e1718	e1819	95% CI	Delta	% Change
all	196.65	234.81	(202.38, 267.24)	38.16	19%

The estimated clearing area for the 2018-19 monitoring period in the Brigalow Belt was 35,550 hectares in remnant areas and 199,270 hectares in non-remnant areas (Table 8). This represents a 58% and 14% increase, respectively, compared to the previous monitoring period. The 95% confidence interval suggests that this difference is significant for remnant clearing. This estimate is consistent with the previous three monitoring periods (Figure 3).

Table 8: Summary of estimated clearing area for the Brigalow Belt bioregion in the 2018-19 period

Veg Class	Area (1000's ha)			Change	
	e1718	e1819	95% CI	Delta	% Change
Remnant	22.46	35.55	(28.10, 42.99)	13.09	58%
Non-remnant	174.19	199.27	(169.65, 228.89)	25.07	14%





Figure 3. Clearing areas for Brigalow Belt bioregion from 2013-14 to 2018-19 for all clearing (top) and separated into clearing in remnant and non-remnant areas (bottom). The sampling-based estimates are presented as grey bars, with the 95% confidence interval superimposed.

6.1.4 Conclusion

The results of this study estimate that using the SLATS Landsat-based methodology, there was 234,810 hectares of clearing in the Brigalow Belt in 2018-19. This would be an increase of 19% in the area cleared in the Brigalow Belt in 2018-19 compared to the 2017-18 monitoring period which had 196,650 hectares of clearing mapped. The estimates of total clearing are consistent with previous recent monitoring periods. Importantly, there is no evidence that clearing rates have reduced between the 2017-18 and 2018-19 monitoring periods.

There are some difficulties in implementing a sampling approach to faithfully recreate the conditions that would have been in place if the SLATS program had continued with Landsat instead of moving to Sentinel-2 data. Significant user variability, for example, was identified in this exercise, and given the small size of the team available to do the sampling exercise could have introduced some bias. It should be noted though that the results also aligned with observations of senior scientists from their Sentinel-2-based mapping for the 2018-19 monitoring

period.

The study demonstrated a method for estimating a clearing area for a given region, based on a stratified sample. Areas estimates were obtained, with confidence intervals. Thus, comparisons could be made with previously reported clearing figures to determine if clearing had significantly increased, decreased, or remained at similar levels as the previous monitoring period(s). The method therefore represents an efficient approach for obtaining clearing estimates if the aim is to provide indicative figures to guide decision-making. The sample-based method is similar to statistical sampling approaches applied in other areas of official reporting, such as is undertaken by the Australian Bureau of Statistics. It is therefore not a new concept in the context of official reporting and could be considered where future comparable estimates of Landsat-based clearing mapping are required. However, it is anticipated that once the Sentinel-2 based program undertakes two or three monitoring periods, this requirement may diminish. That said, the historical context that the SLATS Landsat clearing record provides with respect to environmental and policy change in Queensland is pivotal to understanding the current and future state of Queensland's ecosystems and the policy which is intended to protect them while maintaining sustainable development, particularly for agriculture. The method presented has the potential to continue to contribute to that record.